Structure of Tate-Shafarevich groups of elliptic curves over global function fields

M.L. Brown

Institut Fourier, B.P. 74, 38402 Saint Martin d'Hères, France

arXiv:1602.02932v1 [math.NT] 9 Feb 2016

Shortened Title
Tate-Shafarevich Groups

Abstract

The structure of the Tate-Shafarevich groups of a class of elliptic curves over global function fields is determined. These are known to be finite abelian groups from the monograph [1] and hence they are direct sums of finite cyclic groups where the orders of these cyclic components are invariants of the Tate-Shafarevich group. This decomposition of the Tate-Shafarevich groups into direct sums of finite cyclic groups depends on the behaviour of Drinfeld-Heegner points on these elliptic curves. These are points analogous to Heegner points on elliptic curves over the rational numbers.

AMS 2012 Mathematics Subject Classification Numbers
11G05, 11G09, 11G20, 11G40, 14G10, 14G17, 14G25, 14H52

Keywords: Elliptic curves, Tate-Shafarevich groups, Function fields

Table of Contents

Chapter 1. Preliminaries
- 1.1. Introduction
- 1.2. Global fields of positive characteristic
- 1.3. Orders in imaginary quadratic fields
- 1.4. Ring class fields
- 1.5. Elliptic curves over global fields of positive characteristic
- 1.6. The Drinfeld modular curve $X_0^{\mathrm{Drin}}(I)$
- 1.7. Analogue for F of the Shimura-Taniyama-Weil conjecture
- 1.8. Drinfeld-Heegner points
- 1.9. Groups and cohomology

1.10. Torsion on elliptic curves E/F
1.11. Igusa's theorem
1.12 Consequences of Igusa's theorem
Chapter 2. Local duality, Cassels pairings, Tate-Shafarevich groups
2.1. Local duality of elliptic curves
2.2. Selmer groups and Tate-Shafarevich groups
2.3. The Cassels pairing
Chapter 3. The cohomology classes $\gamma_n(c), \delta_n(c)$
3.1. The set $\mathcal{P}$ of prime numbers
3.2. Frobenius elements and the set $\Lambda(n)$ of divisors
3.3. A refined Hasse principle for finite group schemes
3.4. Drinfeld-Heegner points and the cohomology classes $\gamma_n(c)$, $\delta_n(c)$
Chapter 4. Structure of the Tate-Shafarevich group and the Selmer group
4.1. Statement of the main theorems
4.2. Cochains for the the cohomology classes $\gamma_n(c), \delta_n(c)$
4.3. Points P_c defined over local fields
4.4. The map χ_z
4.5. Localizations of the classes $\gamma_n(c)$ and $\delta_n(c)$
4.6. The Cassels pairing with a class $\delta_n(c)$
Chapter 5. Construction of cohomology classes and proofs of the main theorems
5.1. M_r is finite for some r
5.2. A class $\gamma_n(c)$ in the Selmer group
5.3. Proof of Theorem 4.1.15
5.4. Proof of Theorems 4.1.9 and 4.1.13
5.5. Proof of Theorems 1.1.1 and 4.1.14
5.6. Generators of Tate-Shafarevich groups

Chapter 1. Preliminaries

1.1. Introduction

Let F be a global field of positive characteristic $p > 0$. Let E/F be an elliptic curve with an origin, that is to say a 1-dimensional abelian variety.

In [1] it is shown that for a class of these elliptic curves E/F, the Tate-Shafarevich group $Ш(E/F)$ is finite and for prime numbers l belonging to a set $\mathcal{S}$ of prime numbers given by arithmetic conditions then the l-primary component $Ш(E/F)_{l^\infty}$ has order which is explicitly bounded.

In this paper, we determine the structure of the finite abelian group $Ш(E/F)_{l^\infty}$ for the same class of elliptic curves and for all l in the same set of prime numbers $\mathcal{S}$ (in the notation of Theorem 4.1.9 below $\mathcal{S}$ is the set $\mathcal{P}$ with the exclusion of the prime divisors of the order of the Picard group $\mathrm{Pic}(A)$).

We also determine the structure of the Selmer groups of the elliptic curves in question.

Let E/F be an elliptic curve and let K be an imaginary quadratic extension of F with respect to the place ∞ of F (see §1.2). Let Spec A be the non-singular affine curve with function field F and whose point at infinity is ∞ (see §1.2). Assume that E, K, ∞ satisfy the hypotheses (a), (b), (c) of (4.1.1) in chapter 4. This provides an infinite set of prime numbers $\mathcal{P}$ of positive Dirichlet density and defined by arithmetic conditions (see §3.1). Indeed, $\mathcal{P}$ contains all except finitely many prime numbers $l \in \mathbb{Z}$ of the form $2^s n + 1$ where $s \geq 1$ and n is odd such that q is a 2^sth power non-residue modulo l where q is the order of the exact finite field of constants of F.

Fix a prime number $l \in \mathcal{P}$. There are sets of divisors $\Lambda^r(n)$, relative to l, on F for all integers $n \geq 1$ such that each divisor in $\Lambda^r(n)$ is a sum of r distinct prime divisors and there is a decreasing filtration on $\Lambda^r(1)$.

$$\Lambda^r = \Lambda^r(1) \supseteq \Lambda^r(2) \supseteq \dots$$

For any divisor $c \in \Lambda^r(n)$ there is a corresponding Drinfeld-Heegner point P_c of $E(K[c])$, the group of $K[c]$-rational points of E, where $K[c]$ is the ring class field of K with conductor c (see §1.4 and (3.4.9)).

On $E(K[c])$ there is the decreasing l-adic filtration

$$E(K[c]) \supseteq lE(K[c]) \supseteq l^2E(K[c]) \supseteq \dots$$

Define

$$M_r = \min_{c \in \Lambda^r}(\max(n \in \mathbb{N} \mid P_c \in l^n E(K[c]))) \text{ for all integers } r \in \mathbb{N}.$$

If the point $P_0 \in E(K)$ has infinite order in the group of K-rational points $E(K)$ then it can be shown that $M_0, M_1, \dots$ is a decreasing sequence of non-negative integers (see Lemma 5.1.4). One of the main results of this paper is the following.

1.1.1. Theorem. *Suppose that P_0 has infinite order in $E(K)$, the group of K-rational points of E. Let l be a prime number in $\mathcal{P}$ and which is coprime to the order of the Picard group of the affine curve* Spec *A. Let $\epsilon = \pm 1$ be the sign in the functional equation of the L-function of E/F. Then the Tate-Shafarevich group $Ш(E/F)$ of E/F is finite and its l-primary component is given by*

$$Ш(E/F)_{l^\infty} \cong \prod_{\substack{(-1)^i = \epsilon \\ i \geq 0}} (\mathbb{Z}/l^{M_i - M_{i+1}}\mathbb{Z})^2$$

where the product runs over integers $i \in \mathbb{N}$ such that $(-1)^i = \epsilon$.

A similar statement holds for the Tate-Shafarevich group of the elliptic curve $E \times_F K$ over K (see Theorem 4.1.9) as well as the Selmer groups of these curves (corollary 4.1.19). The main results of this paper are stated in §4.1.

It may be conjectured that for every global field F of characteristic > 0 there are infinitely many non-isomorphic elliptic curves E/F and infinitely many imaginary quadratic field extensions K/F such that E, K, ∞ satisfy the hypotheses of this Theorem 1.1.1 and those of §4.1. If this conjecture holds, then the above theorem and those of §4.1 give infinitely many non-isomorphic elliptic curves over a given global field of positive characteristic whose l-primary components of the Tate-Shafarevich group are structurally known for infinitely many prime numbers l satisfying arithmetic conditions.

The method of this paper is related to that of Kolyvagin's determination of the structure of Tate-Shafarevich groups of a class of elliptic curves over the rational numbers (see [6] and [8]).

The proofs of the main theorems of this paper stated in §4.1 and Theorem 1.1.1 above require many preliminary results which are explained in chapters 1-5.

Chapter 2 contain basics on Tate local duality, Selmer groups, and the Cassels pairing on Tate-Shafarevich groups. In section 3.1, the set $\mathcal{P}$ of prime numbers is defined by arithmetic conditions. In section §3.2, the sets $\Lambda^r(n)$ of divisors on the global field F are defined. Sections §§3.3,3.4 constructs the cohomology classes $\gamma_n(c), \delta_n(c)$ in the cohomology of the elliptic curve E/F.

In section §4.1., the main results of this paper are stated which are then proved in §§5.3-5.5 after some further properties of $\gamma_n(c), \delta_n(c)$ are proved in Chapters 4 and 5. We show in particular that the cohomology classes $\delta_n(c)$ define characters, via the Cassels pairing on $Ш(E/F)_{l^\infty}$ which determine the structure of this group. The method of proof of the main results in §4.1. is by the construction of many independent elements of the Tate-Shafarevich group $Ш(E/F)_{l^\infty}$. Finally, §5.6 contains complements to the main results.

While care has been taken to minimize the number of hypotheses required for the main theorems of this paper, these hypotheses are still numerous (see for example §3.1, Definition 3.1.3). The assiduous reader will have an abundance of interesting problems in their elimination.

1.2. Global fields of positive characteristic

The notation of this paper is mainly that of the monograph [1] and is detailed in the rest of this chapter 1. A few differences arise, notably the sets of divisors $\Lambda(n)$, which are required for the more refined results of this paper.

Let

k be a finite field of characteristic p with $q = p^m$ elements;
$\overline{k}$ be an algebraic closure of k;

C/k be a smooth projective irreducible curve over k;
F be the function field of C; these hypotheses imply that the finite field k is the exact field of constants of the global field F;
Σ_L, for any global field L, be the set of all places of the field L;
$\infty \in \Sigma_F$ be a closed point of C/k;
$\kappa(z)$ be the residue field at a place $z \in \Sigma_F$ of F;
F_v be the completion of F at the place $v \in \Sigma_F$;
F^{sep} be the separable closure of F;
A be the coordinate ring $\Gamma(C \setminus \{\infty\}, \mathcal{O}_C)$ of the affine curve $C \setminus \{\infty\}$;
$\text{Div}_+(A)$ be the semi-group of effective k-rational divisors on Spec A; that is to say $\text{Div}_+(A)$ is the semi-group of effective k-rational divisors on C/k which are coprime to the place ∞; an element of $\text{Div}_+(A)$ may be written as a finite linear combination $\sum_i n_i z_i$ where $n_i \in \mathbb{N}$ and z_i are prime divisors on Spec A for all i;
$\text{Supp}(c)$, for a divisor $c \in \text{Div}_+(A)$, be the support of the divisor c which is the set of prime divisors with non-zero coefficient in c;
$\text{Pic}(A)$ be the Picard group of A, the group of projective A-modules of rank 1;
K be a separable imaginary quadratic extension field of F with respect to ∞, that is to say K is a quadratic extension field of F in which the place ∞ remains inert;
B be the integral closure of A in K;
τ be the non-trivial element of the galois group $\text{Gal}(K/F)$.

1.3. Orders in imaginary quadratic field extensions

Let K/F be the imaginary quadratic field extension with respect to ∞ of §1.2.

An *order* O in K with respect to A is an A-subalgebra of B whose fraction field is equal to K.

There is a bijection between orders O_c of K with respect to A and effective k-rational divisors c in $\text{Div}_+(A)$ and it is given by

$$c \mapsto A + BI(c)$$

where $I(c)$ is the ideal of A cutting out the divisor c. The divisor c is the *conductor* of the order O_c.

[For more details on orders in imaginary quadratic extensions, see [1, Chapter 2, §2.2]

1.4. Ring class fields

Let

O_c be the order of K with respect to A and with conductor c
where $c \in \mathrm{Div}_+(A)$ (see §1.3);
A_v, for each place v of A, be the localisation of A at v;
$\widehat{O}_{c,v}$ be the completion of the semi-local ring $O_c \otimes_A A_v$;
$G_c = K_\infty^* \prod_v \widehat{O}_{c,v}^*$ be the subgroup of the idèle group of the global field K
whose components are the units of $\widehat{O}_{c,v}$ for all places $v \neq \infty$ of F
and K_∞^* for the place $v = \infty$ and where in the product v runs over all
places of F;
$K[c]$, for any divisor $c \in \mathrm{Div}_+(A)$, be the ring class field with conductor c
with respect to ∞; this is the finite abelian extension field of K defined
by the subgroup G_c of the idèle group of K via the reciprocity map;
$G(c/c')$ be the Galois group of the field extension $K[c]/K[c']$ for divisors $c \geq c'$
of $\mathrm{Div}_+(A)$.

We have these properties of the decomposition of primes in ring class fields (for the proofs, see [1, Chapter 2, §2.3.13]):

(a) The primes ramified in $K[c]/K$ are precisely the primes in the support of c.
(b) The extension $K[c]/K$ is split completely at the place of K lying above ∞.
(c) If $z \notin \mathrm{Supp}(c)$ then for any positive integer n, the galois extension $K[c + nz]/K[c]$ is totally ramified at all places of $K[c]$ above z.

[See [1, Chapter 2, §2.3] for more details on ring class fields.]

1.5. Elliptic curves over global fields of positive characteristic

Let

C/k be a smooth projective irredicible curve over k (as in §1.2);
X/k be an elliptic surface over C, that is to say X/k is a smooth
projective irreducible surface, equipped with a morphism $f : X \to C$
which has a section, such that all fibres of f, except a finite number, are
elliptic curves;
E/F be the generic fibre of $f : X \to C$, which is an elliptic curve E over F
equipped with an origin where F is the function field of C.

The conductor of an elliptic curve E/F is an effective k-rational divisor on F

supported only at the places of bad reduction of E and whose multiplicities are defined in terms of the Galois representation of $\mathrm{Gal}(F^{\mathrm{sep}}/F)$ given by E.

[See [1, §§B.11.1-B.11.4 for the definition of the conductor of E/F.]

1.6. The Drinfeld modular curve $X_0^{\mathrm{Drin}}(I)$

Let

I be a non-zero ideal of A.

$X_0^{\mathrm{Drin}}(I)$ be the curve which is the coarse moduli scheme of Drinfeld modules of rank 2 for A equipped with an I-cyclic structure (see [1, Definition 2.4.2, p. 23]); this curve is compactified by a finite number of cusps which correspond to "degenerate" Drinfeld modules.

This curve $X_0^{\mathrm{Drin}}(I)$ is an analogue for the global field F of the classical modular curve $X_0(N)$ which is the coarse moduli scheme of elliptic curves equipped with a cyclic subgroup of order N, where $N \in \mathbb{N}$.

[For more details, see [1, §2.4].]

1.7. Analogue for F of the Shimura-Taniyama-Weil conjecture

Let E/F be an elliptic curve with split (Tate) multiplicative reduction at ∞. Let I be the non-zero ideal of the ring A which is the conductor, without the place at ∞, of the elliptic curve E/F.

According to the work of Drinfeld on the Langlands conjecture, there is a finite surjective morphism of curves over F

$$X_0^{\mathrm{Drin}}(I) \to E.$$

This result is an analogue for the global field F of the Shimura-Taniyama-Weil conjecture proved by Wiles for semi-stable elliptic curves over the rational numbers.

[For more details see [1, §4.7] and [1, Appendix B].]

1.8. Drinfeld-Heegner points

Let K be an imaginary quadratic extension field of F with respect to ∞ (as in §1.2) and let I be a non-zero ideal of A.

Let D be a rank 2 Drinfeld module for A with complex multiplication by an order $\mathcal{O}$ of the field K with respect to A, that is to say $\mathcal{O}$ is a subring of

K which is integral over A. Let Z be an I-cyclic subgroup of D. Then the pair (D, Z) represents a non-cuspidal point of the modular curve $X_0^{\mathrm{Drin}}(I)$. Such points (D, Z) exist if the prime divisors in the support of I split completely in the field extension K/F.

If the quotient Drinfeld module D/Z has the same ring of endomorphisms $\mathcal{O}$ as D then the point (D, Z) on $X_0^{\mathrm{Drin}}(I)$ is called a *Drinfeld-Heegner point*.

If $f : X_0^{\mathrm{Drin}}(I) \to E$ is a finite morphism of curves where E/F is an elliptic curve (see §§1.5,1.6,1.7), then the point $f(D, Z)$ of the elliptic curve E is also called a Drinfeld-Heegner point.

The Drinfeld-Heegner points (D, Z) and $f(D, Z)$ are rational over the ring class field $K[c]$ where c is the conductor of the order $\mathcal{O}$ of K relative to A.

[See [1, §§2.2,2.3] or §3.4 below for more details.]

1.9. Groups and cohomology

If G is a discrete abelian group denote by

G_m the kernel of multiplication by the integer $m \in \mathbb{N}$ on G;
${}_mG$ the cokernel G/mG of multiplication by the integer $m \in \mathbb{N}$;
$|G|$ the order of the group G which is either a positive integer or $+\infty$;
$\mathrm{ord}(g)$ the order of an element $g \in G$ which is the cardinality of the subgroup generated by g;
$\exp(G)$ the exponent of G which is maximum order of an element of G;
$\widehat{G}$ the Pontrjagin dual of G namely the topological group $\mathrm{Hom}(G, \mathbb{Q}/\mathbb{Z})$;

If G is a finite abelian group then $\widehat{G}$ may be identified with the group of 1-dimensional complex characters of G, that is to say the group of homomorphisms $\mathrm{Hom}(G, \mathbb{C}^*)$. A character of a finite abelian group is always assumed to be irreducible.

If F'/F is a galois extension of a global field F and if $z \in \Sigma_F$ is a prime divisor of F unramified in F' we denote by $\mathrm{Frob}(z)$, or $\mathrm{Frob}_{F'/F}(z)$, the conjugacy class in $\mathrm{Gal}(F'/F)$ of Frobenius substitutions associated to z.

If L is a field, we shall write $H^i(L, M)$ for the Galois cohomology group $H^i(\mathrm{Gal}(L^{\mathrm{sep}}/L), M)$, where L^{sep} is the separable closure of L. If L'/L is a finite galois field extension, we write $H^i(L'/L, M)$ for $H^i(\mathrm{Gal}(L'/L), M)$.

If F is a global field and z is a place of F then the restriction, or localization, of a class $c \in H^i(L, M)$ is written $c_z \in H^i(F_z, M)$ where F_z is the completion of F at z.

1.10. Torsion on elliptic curves E/F

Let E be an elliptic curve over a global field F of positive characteristic $p > 0$ as in §1.5. Let K be an imaginary quadratic extension field of F with respect to the place ∞ of F. As in §1.4, let $K[c]$ be the ring class field over K with conductor $c \in \mathrm{Div}_+(A)$. Put

$$K[A] = \bigcup_{c \in \mathrm{Div}_+(A)} K[c].$$

That is to say $K[A]$ is a field which is the union of all the ring class fields $K[c]$ in some algebraic closure of K.

Let S be a subset of $\mathbb{N}^*$ such that if $n_1 \in S$ and n_0 is any divisor of n_1 then $n_0 \in S$. A *quasi-group* $\{G_n\}_{n \in S}$ relative to S is a family of abelian groups G_n indexed by the elements n of S such that $nG_n = 0$ and if $n_0, n_1 \in S$ are elements where n_0 divides n_1 there is a group homomorphism $f_{n_1 n_0} : G_{n_1} \to G_{n_0}$ satisfying the compatibility condition that if $n_0, n_1, n_2 \in S$ and n_0 divides n_1 and n_1 divides n_2 then $f_{n_2 n_0} = f_{n_1 n_0} \circ f_{n_2 n_1}$. [See [1, Chapter 7, §7.1, p. 330].]

1.10.1 Proposition. ([1, Proposition 7.3.8.]) *The quasi-group*

$$\{E(K[A])_n\}_{n \in \mathbb{N}}$$

is trivial, that is to say the order of the torsion group $E(K[A])_n$ is bounded independently of n and there is a finite set $\mathcal{E}$ of prime numbers such that for all integers n prime to all elements of $\mathcal{E}$ we have

$$E(K[A])_n \cong 0. \quad \square$$

1.10.2. Proposition. ([1, Proposition 7.14.2]) *Let $\mathcal{E}$ be the finite set of prime numbers of Proposition 1.10.1. For any divisor c of $\mathrm{Div}_+(A)$, the restriction homomorphism*

$$H^1(K, E_n) \to H^1(K[c], E_n)^{\mathcal{G}_c}$$

is an isomorphism for all integers n prime to $\mathcal{E}$ where $\mathcal{G}_c = \mathrm{Gal}(K[c]/K)$.

This follows from Proposition 1.10.1 and the Hochschild-Serre spectral sequence

$$H^i(\mathcal{G}_c, H^j(K[c], E_n(K^{\mathrm{sep}}))) \Rightarrow H^{i+j}(K, E_n(K^{\mathrm{sep}}))$$

(more details are given in [1, Proposition 7.14.2]).□

1.11. Igusa's theorem

This section is a summary of the results of Igusa for the Galois action on torsion points of elliptic curves over global fields of positive characteristic.

(1.11.1) As in §1.2, let F be a global field of positive characteristic p where k is the exact field of constants of F and let E/F be an elliptic curve. Let

$G = \mathrm{Gal}(F^{\mathrm{sep}}/F)$, where F^{sep} is the separable closure of F;
n be an integer prime to p;
E_n be the finite F-group scheme of n-torsion points of E;
E_∞ be the torsion subgroup of $E(F^{\mathrm{sep}})$ of order prime to p.

The elliptic curve E/F is said to be *isotrivial* if there is a finite galois extension field F' of F such that the curve $E \times_F F'$ is definable over a finite subfield of F'.

(1.11.2) The action of the galois group G on E_n provides a homomophism

$$\rho_n : G \to \mathrm{Aut}(E_n) \cong \mathrm{GL}_2(\mathbb{Z}/n\mathbb{Z}).$$

The determinant

$$\det : \mathrm{Aut}(E_n) \to (\mathbb{Z}/n\mathbb{Z})^*$$

induces a homomorphism

$$G \to (\mathbb{Z}/n\mathbb{Z})^*.$$

Let H_n be the subgroup of $(\mathbb{Z}/n\mathbb{Z})^*$ generated by the powers of $q = |k|$ modulo n. Then H_n is naturally isomorphic to the Galois group of the field of nth roots of unity over k. Let Γ_n be the subgroup of $\mathrm{GL}_2(\mathbb{Z}/n\mathbb{Z})$ defined by the exact sequence of finite groups, where det is the restriction to Γ_n of the determinant homomorphism on $\mathrm{GL}_2(\mathbb{Z}/n\mathbb{Z})$,

(1.11.3)
$$0 \to \mathrm{SL}_2(\mathbb{Z}/n\mathbb{Z}) \to \Gamma_n \xrightarrow{\det} H_n \to 0.$$

(1.11.4) Passing to the projective limit of the previous exact sequence over all integers n prime to p we obtain the exact sequence of profinite groups

$$0 \longrightarrow \mathrm{SL}_2(\widehat{\mathbb{Z}}^{(p)}) \longrightarrow \widehat{\Gamma} \longrightarrow \widehat{H} \longrightarrow 0$$

where $\widehat{H}$ is the subgroup of $\widehat{\mathbb{Z}}^{(p)*}$ topologically generated by q, where

$$\widehat{\mathbb{Z}}^{(p)} = \prod_{l \neq p} \mathbb{Z}_l$$

is the profinite prime-to-p completion of $\mathbb{Z}$, and $\widehat{\Gamma}$ is a closed subgroup of $\mathrm{GL}_2(\widehat{\mathbb{Z}}^{(p)})$.

(1.11.5) Passing to the projective limit of the exact sequence (1.11.3) where n runs over all powers of a prime number l where $l \neq p$, we obtain the exact sequence

$$0 \longrightarrow \mathrm{SL}_2(\mathbb{Z}_l) \longrightarrow \widehat{\Gamma}_l \longrightarrow \widehat{H}_l \longrightarrow 0.$$

1.11.6. Theorem. (Igusa [I]). *Suppose that E/F is not isotrivial. Then the profinite group* $\mathrm{Gal}(F(E_\infty)/F)$ *is an open subgroup of* $\widehat{\Gamma}$. $\square$

This result has the following consequence.

1.11.7. Theorem. *Suppose that E/F is not isotrivial. Then for all prime numbers $l \neq p$ the profinite group* $\mathrm{Gal}(F(E_{l^\infty})/F)$ *is an open subgroup of* $\widehat{\Gamma}_l$ *and is equal to* $\widehat{\Gamma}_l$ *for all but finitely many l.* $\square$

1.11.8. Remarks. (1) Suppose that the curve E/F is isotrivial. Then it is easy to show that the group $\mathrm{Gal}(F(E_\infty)/F)$ is an extension of a finite group by the abelian profinite group $\widehat{\mathbb{Z}}^{(p)}$.
(2) Let E be an elliptic curve defined over a number field L. The galois action on the torsion points of E/L is known and depends principally on whether or not E has complex multiplication.

[See [S5] and [S5, §4.5] for both the cases of complex multiplication and without complex multiplication. See also [1, Chapter 7, Remarks 7.2.8] for more details.]

1.12. Consequences of Igusa's theorem

For a finite group G and a $\mathbb{Z}[G]$-module M, let $H^i(G, M)$ denote the standard cohomology groups of G acting on M (see [1, §5.6], see also [9, Chap. 1] for the Tate modified cohomology groups).

1.12.1. Proposition. *Let E/F be an elliptic curve and let $\mathbb{N}^{(p)}$ be the set of positive integers coprime to p, where p is the characteristic of F. Write G_n for the group* $\mathrm{Gal}(F(E_n)/F)$.
(i) *Let $i = 0$ or 1. Then*

$$\{H^i(G_n, E_n)\}_{n \in \mathbb{N}^{(p)}}$$

is a trivial quasi-group.
(ii) *There is a finite set $\mathcal{N}$ of prime numbers including p such that for all prime numbers $l \notin \mathcal{N}$ we have*

$$H^i(G_{l^n}, E_{l^n}) = 0 \quad \textit{for all} \quad n \geq 1 \quad \textit{and all} \quad i \geq 0.$$

[Part (i) may be restated as: for $i = 0$ or 1 and for all n coprime to p, the order of the group $H^i(G_n, E_n)$ is bounded independently of n and there is a finite set of prime numbers such that for all integers n coprime to this set of prime numbers we have $H^i(G_n, E_n) \cong 0$. For the proof of this proposition, see [1, Chapter 7, Proposition 7.3.1] $\square$

(1.12.2) For each prime number l different from p, select once and for all a basis of the Tate module $T_l(E)$ over $\mathbb{Z}_l$, the l-adic completion of $\mathbb{Z}$; this fixes for the rest of this paper, for every prime number $l \neq p$, an isomorphism of $\mathrm{Gal}(F(E_{l^\infty})/F)$ with a subgroup of $\mathrm{GL}_2(\mathbb{Z}_l)$ and the two groups may then be identified with each other.

1.12.3. Proposition. *Let E/F be an elliptic curve which is not isotrivial. Let L be a finite extension field of F in which k is algebraically closed. Then there is an infinite set S of prime numbers of positive Dirichlet density such that for all $l \in S$ we have*
(a) *the fields $F(E_{l^\infty})$ and L are linearly disjoint over F;*
(b) *$E(L)_{l^\infty} = 0$;*
(c) $\begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} \in \mathrm{Gal}(F(E_{l^\infty})/F)$.

[For the proof, see [1, Proposition 7.3.10].] □

Chapter 2. Local duality, Cassels pairings, Tate-Shafarevich groups

2.1. Local duality of elliptic curves

This section is a brief summary of Tate-Poitou local duality for elliptic curves over a local field.

[For more details on local duality of abelian varieties, see [9, Chapter I and Chapter III, §7].]

(2.1.1) Let

L be a non-archimedian complete local field;
L^{sep} be the separable closure of L;
E/L be an elliptic curve over L;
$n \geq 1$ be an integer coprime to the characteristic of L;
G be the Galois group $\mathrm{Gal}(L^{\mathrm{sep}}/L)$.

(2.1.2) Let μ_n be the multiplicative subgroup of L^{sep} of nth roots of unity. Then μ_n is a finite G-module. Let E_n denote the G-module of n-torsion points of $E(L^{\mathrm{sep}})$. We have an abelian group isomorphism

$$E_n \cong \frac{\mathbb{Z}}{n\mathbb{Z}} \oplus \frac{\mathbb{Z}}{n\mathbb{Z}}.$$

Denote by $\{,\}$ the Weil pairing

$$\{,\} : E_n \times E_n \to \mu_n.$$

This is a perfect pairing of G-modules. In particular, we have an isomorphism of G-modules

$$E_n \cong \mathrm{Hom}_G(E_n, \mu_n).$$

(2.1.3) The Weil pairing induces a cup-product pairing in Galois cohomology

$$H^1(L, E_n) \times H^1(L, E_n) \to H^2(L, \mu_n).$$

This is a non-degenerate anti-symmetric pairing of abelian groups. By local class field theory, we have a canonical isomorphism of groups, where $\mathrm{Br}(L)$ is the Brauer group of L,

$$\mathrm{Br}(L) \cong \frac{\mathbb{Q}}{\mathbb{Z}}.$$

This induces an isomorphism

$$H^2(L, \mu_n) = \mathrm{Br}(L)_n \cong \frac{\mathbb{Z}}{n\mathbb{Z}}.$$

2.1.4. Theorem. (Tate-Poitou local duality). *The cup product pairing*

$$<,>_v: H^1(L,E_n)\times H^1(L,E_n)\to \frac{\mathbb{Z}}{n\mathbb{Z}} \tag{2.1.5}$$

obtained from the Weil pairing is an alternating and non-degenerate pairing of $\frac{\mathbb{Z}}{n\mathbb{Z}}$*-modules.*

[For the proof, see [9, Chapter I, Cor. 2.3].]□

2.1.6. Theorem. *Assume that* n *is prime to the residue field characteristic of* L*.*
(i) *The subgroup* ${}_nE(L)$ *of* $H^1(L,E_n)$ *is isotropic for the alternating pairing* $<,>_v$*.*
(ii) *The cup product pairing* $<,>_v$ *on* $H^1(L,E_n)$ *induces a non-degenerate pairing of abelian groups, where* ${}_nE(L)=E(L)/nE(L)$*,*

$$[,]_v : {}_nE(L)\times H^1(L,E)_n\to \frac{\mathbb{Z}}{n\mathbb{Z}}.$$

[For the proof, see [1, Chapter 7, Theorem 7.15.6, p.403] for part (i) and [9, Chap. I, Cor. 3.4 and Remark 3.6, Chapter III, Theorem 7.8] for part (ii). Note that part (ii) holds without the hypothesis that n be coprime to the characteristic of L, see [9, Chap. III, §7].] □

(2.1.7) Suppose now that K is a global field of positive characteristic, Σ_K is the set of all places of K, and that the integer n is coprime to the characteristic of K. Let E/K be an elliptic curve.

2.1.8. Proposition. *Let* c *and* c' *be two elements of* $H^1(K,E_n)$*. Denote by* c_v *and* c'_v *the induced elements of* $H^1(K_v,E_n)$ *for all places* $v\in\Sigma_K$ *of* K *where* K_v *is the completion of* K *at* v*. Then we have*

$$\sum_{v\in\Sigma_K} < c_v, c'_v >_v = 0.$$

Proof. The sum of the local invariants of a global class in $H^2(K,\mathbb{G}_m)$ is zero.□

2.2. Selmer groups and Tate-Shafarevich groups

(2.2.1) Let

E/F be an elliptic curve as in §1.5;
$n \in \mathbb{N}$ be an integer coprime to the characteristic p of the global field F.

(2.2.2) For a place v of the field F, we write F_v for the completion of F at the place v (as in §1.2). The exact sequence of commutative group schemes over F, obtained from the morphism of multiplication by n,

$$0 \longrightarrow E_n \longrightarrow E \xrightarrow{n} E \longrightarrow 0$$

gives rise to a commutative diagram, where the maps res_v are the restriction homomorphisms at v and the rows are exact sequences of abelian groups,

$$\begin{array}{ccccccccc} 0 & \to & {}_nE(F) & \to & H^1(F, E(F^{\mathrm{sep}})_n) & \to & H^1(F, E(F^{\mathrm{sep}}))_n & \to & 0 \\ & & \downarrow \mathrm{res}_v & & \downarrow \mathrm{res}_v & & \downarrow \mathrm{res}_v & & \\ 0 & \to & {}_nE(F_v) & \to & H^1(F_v, E(F_v^{\mathrm{sep}})_n) & \to & H^1(F_v, E(F_v^{\mathrm{sep}}))_n & \to & 0 \end{array}$$

As in §1.9, ${}_nE(F)$ denotes the cokernel $E(F)/nE(F)$ and $E(F^{\mathrm{sep}})_n$ denotes the n-torsion subgroup of $E(F^{\mathrm{sep}})$.

(2.2.3) The *Tate-Shafarevich group* $Ш(E/F)$ of E/F is defined as

$$Ш(E/F) = \ker\{H^1(F, E) \to \prod_{v \in \Sigma_F} H^1(F_v, E)\}.$$

This group $Ш(E/F)$ is known to be a torsion of cofinite type (see [10]).

The *n-Selmer group* is defined as, in terms of the commutative diagram of (2.2.2) and where res_v denotes the middle vertical homomorphism of the diagram,

$$\mathrm{Sel}_n(E/F) = \bigcap_{v \in \Sigma_F} \mathrm{res}_v^{-1}({}_nE(F_v)).$$

Therefore $\mathrm{Sel}_n(E/F)$ is a subgroup of $H^1(F, E(F^{\mathrm{sep}})_n)$ and is a finite abelian group. We then have the exact sequence of torsion abelian groups from the commutative diagram of (2.2.2), where $Ш(E/F)_n$ is the n-torsion subgroup of $Ш(E/F)$,

$$0 \to {}_nE(F) \to \mathrm{Sel}_n(E/F) \to Ш(E/F)_n \to 0.$$

(2.2.4) Let F'/F be a finite separable galois field extension. We write $Ш(E/F')$ in place of $Ш(E \times_F F'/F')$ for the Tate-Shafarevich group of $E \times_F F'$ over F' obtained by ground field extension from F to F'; similarly, for the Selmer quasi-group, we write $\mathrm{Sel}_n(E/F')$ in place of $\mathrm{Sel}_n(E \times_F F'/F')$.

2.2.5. Proposition. *Let r be the degree of the finite separable field extension F'/F. For any torsion abelian group $\mathcal{A}$ write $\mathcal{A}_{(\text{non } \triangledown)}$ for the torsion subgroup of $\mathcal{A}$ of order coprime to r. The restriction homomorphism provides isomorphisms for all integers n coprime to r*

$$Ш(E/F)_{(\text{non } r)} \overset{\text{res}}{\cong} Ш(E/F')_{(\text{non } r)}{}^{\mathrm{Gal}(F'/F)}$$

$$\mathrm{Sel}_n(E/F) \overset{\text{res}}{\cong} \mathrm{Sel}_n(E/F')^{\mathrm{Gal}(F'/F)}.$$

Proof. The definition of the Tate-Shafarevich groups provides a commutative diagram with exact rows

$$\begin{array}{ccccccc}
0 & \to & Ш(E/F')^{\mathrm{Gal}(F'/F)} & \to & H^1(F',E)^{\mathrm{Gal}(F'/F)} & \to & \Big(\prod_{v\in\Sigma_{F'}} H^1(F'_v,E)\Big)^{\mathrm{Gal}(F'/F)} \\
 & & \uparrow & & \uparrow & & \uparrow \\
0 & \to & Ш(E/F) & \to & H^1(F,E) & \to & \prod_{v\in\Sigma_F} H^1(F_v,E)
\end{array}$$

For any place v of F, the F_v-algebra $F_v \otimes_F F'$ is étale and is the product of the completions of F' at the places lying over v. The inflation restriction sequence provides isomorphisms for any integer s coprime to the order of $\mathrm{Gal}(F'/F)$

$$\begin{aligned}
H^1(F,E)_s &\cong H^1(F',E)_s^{\mathrm{Gal}(F'/F)} \\
H^1(F_v,E)_s &\cong H^1(F_v\otimes_F F',E)_s^{\mathrm{Gal}(F'/F)} \text{ for all places } v \text{ of } F.
\end{aligned}$$

The isomorphism of the proposition for the Tate-Shafarevich groups now follows by a diagram chase. The corresponding isomorphism for the n-Selmer groups follows from the commutative diagram with exact rows

$$\begin{array}{ccccccccc}
0 & \to & {}_nE(F) & \to & \mathrm{Sel}_n(E/F) & \to & Ш(E/F)_n & \to & 0 \\
 & & \downarrow\cong & & \downarrow & & \downarrow\cong & & \\
0 & \to & ({}_nE(F'))^{\mathrm{Gal}(F'/F)} & \to & \mathrm{Sel}_n(E/F')^{\mathrm{Gal}(F'/F)} & \to & Ш(E/F')_n^{\mathrm{Gal}(F'/F)} & \to & 0
\end{array}$$

as required. □

2.2.6. Remark. This section is a generalised form of [1, Chapter 7, §7.9].

2.3. The Cassels pairing

(2.3.1) Let E/F be an elliptic curve over the global field F of characteristic $p > 0$ as in §1.5. The Tate-Shafarevich group $Ш(E/F)$ of E/F is equipped with the anti-symmetric Cassels pairing

$$<,>_{\text{Cassels}} : Ш(E/F) \times Ш(E/F) \to \mathbb{Q}/\mathbb{Z}$$

which is non-degenerate if $Ш(E/F)$ is finite. The Tate-Shafarevich group $Ш(E/F)$ is a torsion group of cofinite type (see [10]).

In this section, the Cassels pairing is defined for E/F for the non-p part $Ш(E/F)_{(\text{non}-p)}$ of $Ш(E/F)$, that is to say the subgroup of the Tate-Shafarevich group of order coprime to the characteristic p.

(2.3.2) For any place $v \in \Sigma_F$ of F, we have the commutative diagram with exact rows obtained from restriction from F to F_v for any integer m where ∂_m is the connecting homomorphism induced from the morphism of multiplication by m on E

$$\begin{array}{ccccccccccc}
0 & \to & E(F)_m & \to & E(F) & \to & E(F) & \stackrel{\partial_m}{\to} & H^1(F, E_m) & \to & H^1(F, E)_m \\
 & & \downarrow & & \downarrow & & \downarrow & & \downarrow & & \downarrow \\
0 & \to & E(F_v)_m & \to & E(F_v) & \to & E(F_v) & \stackrel{\partial_m}{\to} & H^1(F_v, E_m) & \to & H^1(F_v, E)_m
\end{array}$$

(2.3.3) Let $a, b \in Ш(E/F)_{(\text{non}-p)}$. Let $m \geq 1$ be the order of a and $n \geq 1$ be the order of b where m, n are coprime to the characteristic p of F. Then we have

$$a \in Ш(E/F)_m \text{ and } b \in Ш(E/F)_n.$$

Select elements

$$a^{(1)} \in H^1(F, E_m) \text{ and } b^{(1)} \in H^1(F, E_n)$$

mapping to a and b respectively in the commutative diagram of (2.3.2).

For any element $h \in H^1(F, E)$ denote by h_v the restriction of h to $H^1(F_v, E)$ for any place v of F and similarly for cochains.

(2.3.4) Suppose first that a is divisible by n in $H^1(F, E)$, say $a = na_1$ where $a_1 \in H^1(F, E)_{mn}$. We may select local points $y_v \in {}_nE(F_v)$ such that

$$\partial_n(y_v) = b_v^{(1)}, \text{ for all places } v \in \Sigma_F,$$

as $b_v^{(1)}$ maps to zero in $H^1(F_v, E)$. Let $a_{1,v}$ denote the localization in $H^1(F_v, E)_{mn}$ of a_1. Note that since $a \in Ш(E/F)_m$ we have $a_{1,v} \in H^1(F_v, E)_n$ for all v. For

any $v \in \Sigma_F$, denote by

$$[,]_v : H^1(F_v, E)_n \times {}_nE(F_v) \to \frac{\mathbb{Z}}{n\mathbb{Z}}.$$

the local pairing as in Theorem 2.1.6. Then the Cassels pairing is given in terms of the local pairing by the formula

$$< a, b >_{\text{Cassels}} = \sum_{v \in \Sigma_F} [a_{1,v}, y_v]_v \tag{2.3.5}$$

where the sum runs over all places of F.

(2.3.6) We have

$$a_{1,v} = i_*(c_{1,v})$$

for some $c_{1,v} \in H^1(F_v, E_n)$ for all v where i is the inclusion of group schemes $E_n \hookrightarrow E$. We then have that the Cassels pairing is also given in terms of the cup product pairing $<,>_v$ of Theorem 2.1.4 by

$$< a, b >_{\text{Cassels}} = \sum_v < c_{1,v}, b_v^{(1)} >_v \tag{2.3.7}$$

as we have for all places v

$$[a_{1,v}, y_v]_v = < c_{1,v}, \partial_n(y_v) >_v .$$

(2.3.8) In the previous two paragraphs, the global element $a_1 \in H^1(F, E)_{mn}$ such that $na_1 = a$ may not exist. Nevertheless, in a suitable sense it always exists locally and in general the Cassels pairing is defined as follows.

Select elements $a^{(1)}$ and $b^{(1)}$ of $H^1(F, E_m)$ and $H^1(F, E_n)$ mapping to a and b respectively. For each valuation v of F, let $a_v^{(1)}$ be the localisation in $H^1(F_v, E_m)$ of $a^{(1)} \in H^1(F, E_m)$. For each valuation v of F, a maps to zero in $H^1(F_v, E)$ and hence $a_v^{(1)}$ lies in the image of $E(F_v)$ under ∂_m. Then we can lift, by the diagram where id is the identity map,

$$\begin{array}{ccc} E(F_v) & \stackrel{\partial_m}{\to} & H^1(F_v, E_m) \\ \text{id} \uparrow & & \uparrow n \\ E(F_v) & \stackrel{\partial_{mn}}{\to} & H^1(F_v, E_{mn}) \end{array}$$

$a_v^{(1)} \in H^1(F_v, E_m)$ to an element $a_{v,1}^{(1)} \in H^1(F_v, E_{mn})$ so that $na_{v,1}^{(1)} = a_v^{(1)}$ and $a_{v,1}^{(1)}$ is in the image of $E(F_v)$ under ∂_{mn}.

Let β be a cocycle in $\text{Cocy}^1(F, E_m)$ representing $a^{(1)} \in H^1(F, E_m)$ and lift it to a cochain $\beta_1 \in \text{Coch}^1(F, E_{mn})$. Select a cocycle $\beta_{v,1} \in \text{Cocy}^1(F_v, E_{mn})$ representing the element $a_{v,1}^{(1)} \in H^1(F_v, E_{mn})$ and a cocycle $\beta' \in \text{Cocy}^1(F, E_n)$ representing $b^{(1)} \in H^1(F, E_n)$.

The coboundary $d\beta_1$ of β_1 takes values in E_n as β_1 is a cochain lifting the cocycle β with values in E_m. The cup product $d\beta_1 \cup \beta'$ represents an element of $H^3(\mathrm{Gal}(F^{\mathrm{sep}}/F), \mathbb{G}_m)$ where $\mathbb{G}_m$ is the multiplicative group scheme over F and where this cup product is induced by the Weil pairing $E_n \times E_n \to \mathbb{G}_m$ (see (2.1.3)). But this last cohomology group $H^3(\mathrm{Gal}(F^{\mathrm{sep}}/F), \mathbb{G}_m)$ is zero (see [9, Chap. 1, 4.18 or 4.21]). Hence we have

$$d\beta_1 \cup \beta' = d\epsilon$$

for some 2-cochain $\epsilon \in \mathrm{Coch}^2(F, \mathbb{G}_{\mathbf{m}})$.

The cochain localized at v

$$((\beta_{v,1} - \beta_{1,v}) \cup \beta'_v) + \epsilon_v$$

is a 2-cocycle in $\mathrm{Cocy}^2(F_v, \mathbb{G}_m)$. Denote by

$$\mathrm{inv}_v : \mathrm{Br}(F_v) = H^2(F_v, \mathbb{G}_m) \to \mathbb{Q}/\mathbb{Z}$$

the canonical isomorphism (the invariant map) given by local class field theory. Define

$$< a, b >_{\mathrm{Cassels}} = \sum_{v \in \Sigma_F} \mathrm{inv}_v(((\beta_{v,1} - \beta_{1,v}) \cup \beta'_v) + \epsilon_v) \in \mathbb{Q}/\mathbb{Z}.$$

This can be checked to be independent of the selections made and defines the pairing on $Ш(E/F)_{(\mathrm{non}-p)}$.

2.3.9. Remarks. (1) For this paper, only that part of the construction of the Cassels pairing in paragraphs (2.3.2)-(2.3.6) is required. This is because, under the hypotheses of the main theorems of this paper stated in §4.1, the subgroup $Ш(E/K)$ of $H^1(K, E)$ has the following divisibility property.

Under the hypotheses and notations of the main theorems of this paper stated in §4.1, for all prime numbers $l \in \mathcal{P}$ where l is coprime to $\mathrm{Pic}(A)$ and for any integer $n \geq 0$ there is a subgroup H of $H^1(K, E)$ such that $l^n H = Ш(E/K)_{l^\infty}$.

This divisibility holds because for any sufficiently large positive integer a the group $Ш(E/K)_{l^\infty}$ is generated by the cohomology classes $\delta_{M_0}(c)$ where c ranges of the elements of $\Lambda(a)$ (Theorem 5.6.2) and that $l^n \delta_{M_0+n}(c) = \delta_{M_0}(c)$ if $c \in \Lambda(M_0 + n)$ (lemma 4.2.1(iii)).

It would be interesting to have examples of elliptic curves E/K and prime numbers l for which $Ш(E/K)_{l^\infty}$ is non-zero but does not have this divisibility property as a subgroup of $H^1(K, E)$.

(2) The Cassels pairing for abelian varieties over global fields can be defined in several ways. For a more geometric construction of the pairing than that given above, see [9, Chap. I, Remark 6.11, p. 98]. For the special case of Jacobians of curves, see [9, Chap. I, Remark 6.12, p. 100]. For the construction of the pairing including the p-torsion part of the Tate-Shafarevich group, where p is the characteristic of the base field, see [9, Chap. II, Theorem 5.6, pp. 247-248].

Chapter 3. The cohomology classes $\gamma_n(c), \delta_n(c)$

3.1. The set $\mathcal{P}$ of prime numbers

We define a set $\mathcal{P}$ of prime numbers by arithmetic conditions. For prime numbers l of $\mathcal{P}$ we shall consider in Chapters 4 and 5 the structure of the l-primary component of the Tate-Shafarevich group $Ш(E/F)$ of elliptic curves E/F.

(3.1.1) For a place $v \in \Sigma_F$ of the global field F, let

F_v denote the completion of F at v;
F_v^{nr} denote the maximal unramified extension of the local field F_v
(that is, F_v^{nr} is the field of fractions of the strict henselisation of the valuation ring of F_v);
O_v denote the discrete valuation ring of the local field F_v;
$\infty \in \Sigma_F$ be a place of F;
K be a separable imaginary quadratic extension field of F with respect to the place ∞ as in §1.2;
E/F be an elliptic curve (as in §1.5);
$\mathcal{E}$ denote the Néron model over O_v of the elliptic curve $E \times_F F_v/F_v$;
$\mathcal{E}_0$ denote the closed fibre of $\mathcal{E}/O_v$;
$\pi_0(\mathcal{E}_0)$ be the group of connected components of $\mathcal{E}_0$
as a $\mathrm{Gal}(F_v^{\mathrm{nr}}/F_v)$-module.

Define similarly K_w, K_w^{nr} for a place $w \in \Sigma_K$ of the imaginary quadratic extension field K of F.

3.1.2. Theorem. ([9, Chap. I, Prop. 3.8, p.57]). *Write* $G = \mathrm{Gal}(F_v^{\mathrm{nr}}/F_v)$. *There is an isomorphism*

$$H^1(G, E(F_v^{\mathrm{nr}})) \cong H^1(G, \pi_0(\mathcal{E}_0)).$$

In particular, $H^1(G, E(F_v^{\mathrm{nr}}))$ *is a finite group for all* v *and if the elliptic curve* E *has good reduction at* v *then* $H^1(G, E(F_v^{\mathrm{nr}})) = 0$. □

3.1.3. Definition. Let $\mathcal{P}$ be the set of all prime numbers such that for all $l \in \mathcal{P}$ we have

(a) p, 2, and the prime factors of $|B^*|/|A^*|$ are not in $\mathcal{P}$;
(b) $H^i(K(E_{l^n})/K, E_{l^n}) = 0$ for all integers $n \geq 1$ and for all $i \geq 0$
(see Proposition 1.12.1);

(c) the natural injection $\mathrm{Gal}(F(E_{l^\infty})/F) \to \widehat{\Gamma}_l$ is an isomorphism
(see §1.11 and Igusa's Theorem 1.11.7);
(d) $H^1(K_z^{\mathrm{nr}}/K_z, E)_{l^\infty} = 0$ for all places z of K (see Theorem 3.1.2 above);
(e) K and $F(E_{l^\infty})$ are linearly disjoint over F (see Proposition 1.12.3 or [1, Chapter 7, 7.3.10]);
(f) $\mathcal{P}$ excludes the prime numbers of the finite set $\mathcal{E}$ of Proposition 1.10.1, that is
to say, we have $E(K[A])_{l^m} = 0$ for all $l \in \mathcal{P}$, for all $m \geq 1$, where $K[A]$ is defined in §1.10;
(g) $\begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} \in \mathrm{Gal}(F(E_{l^\infty})/F)$ (see Proposition 1.12.3, which is a consequence of Igusa's Theorem 1.11.6);

3.1.4. Remarks. (1) The first 6 conditions (a),(b),(c),(d),(e),(f) of this definition hold for all except finitely many prime numbers l. Only the last condition (g) fails to hold in general for all but finitely many prime numbers. The set $\mathcal{P}$ is infinite and has positive Dirichlet density.

The set $\mathcal{P}$ can therefore be obtained from the set S of prime numbers provided by Proposition 1.12.3 by deleting a finite number of elements. More precisely, (see [1, Chapter 7, Remarks 7.3.14(1)] and also [1, Chapter 7, Remarks 7.7.6(1)]) the set $\mathcal{P}$ consists of all but finitely many prime numbers l of the form $2^s n + 1$ where $s \geq 1$ and n is odd such that $q = |k|$ is a 2^sth power non-residue modulo l.

(2) The set $\mathcal{P}$ of prime numbers of Definition 3.1.3 above coincides with the set of prime numbers written $\mathcal{P} \setminus \mathcal{F}$ of [1, Lemma 7.14.11]. The only difference between the Definition 3.1.3 above of the set of prime numbers $\mathcal{P}$ and the similar definition [1, Definition 7.10.3] is the extra hypothesis (f) of 3.1.3 above which excludes from $\mathcal{P}$ the finitely many prime numbers of the exceptional set $\mathcal{E}$ of Proposition 1.10.1.

3.1.5. Lemma. *For any integers $0 \leq m \leq n$ and for all prime numbers $l \in \mathcal{P}$ the inclusion of group schemes $E_{l^m} \to E_{l^n}$ induces an injection of cohomology groups*

$$H^1(K, E_{l^m}) \longrightarrow H^1(K, E_{l^n}).$$

Proof of Lemma 3.1.5. This follows from the long exact sequence induced by the isogeny on the finite group scheme E_n of multiplication by l^m

$$0 \to E_{l^m}(K) \to E_{l^n}(K) \to E_{l^{n-m}}(K) \to$$
$$H^1(K, E_{l^m}) \to H^1(K, E_{l^n}) \to H^1(K, E_{l^{n-m}}) \to \dots$$

together with the non-existence of K-rational l-torsion on E (by Proposition 1.10.1 and the definition of $\mathcal{P}$, Definition 3.1.3(f)). □

(3.1.6) We write, where the limits are set-theoretic unions by the previous Lemma 3.1.5,

$$H^1(K, E_{l^\infty}) = \varinjlim_n H^1(K, E_{l^n})$$

and

$$\mathrm{Sel}_{l^\infty}(E/K) = \varinjlim_n \mathrm{Sel}_{l^n}(E/K).$$

(3.1.7) Let $\mathbb{Q}(l)$, for any prime number l, be the additive group of rational numbers with denominators a power of l; the quotient group $\mathbb{Q}(l)/\mathbb{Z}$ is a divisible abelian group where every element is annihilated by a power of l.

We have the exact sequence of abelian groups for all prime numbers $l \in \mathcal{P}$ where $E(K)_{\mathrm{tors}}$ denotes the torsion subgroup of $E(K)$

$$0 \longrightarrow \frac{E(K)}{E(K)_{\mathrm{tors}}} \otimes_{\mathbb{Z}} \frac{\mathbb{Q}(l)}{\mathbb{Z}} \longrightarrow \mathrm{Sel}_{l^\infty}(E/K) \longrightarrow Ш(E/K)_{l^\infty} \longrightarrow 0 \tag{3.1.8}$$

This exact sequence (3.1.8) is obtained from Lemma 3.1.5, the exact sequence of paragraph (2.2.3), and because $E(K)$ has no l-torsion (Definition 3.1.3(f)).

The exact sequence (3.1.8) splits and gives the isomorphism

$$\mathrm{Sel}_{l^\infty}(E/K) \cong \frac{E(K)}{E(K)_{\mathrm{tors}}} \otimes_{\mathbb{Z}} \frac{\mathbb{Q}(l)}{\mathbb{Z}} \quad \oplus \quad Ш(E/K)_{l^\infty}. \tag{3.1.9}$$

This holds as the abelian group $\frac{E(K)}{E(K)_{\mathrm{tors}}} \otimes_{\mathbb{Z}} \frac{\mathbb{Q}(l)}{\mathbb{Z}}$, where $E(K)$ is a finitely generated group, is injective in the category of abelian groups.

It follows from the isomorphism (3.1.9) that $\mathrm{Sel}_{l^m}(E/K)$ is precisely the subgroup of $\mathrm{Sel}_{l^\infty}(E/K)$ annihilated by l^m that is to say we have for all $m \geq 0$ and all $l \in \mathcal{P}$

$$\mathrm{Sel}_{l^m}(E/K) \cong (\mathrm{Sel}_{l^\infty}(E/K))_{l^m}. \tag{3.1.10}$$

3.2. Frobenius elements and the set $\Lambda(n)$ of divisors

(3.2.1) Let

F be the function field of the curve C/k as in §1.2;
$\infty \in \Sigma_F$ be a closed point of C/k;
K be a separable imaginary quadratic extension field of F with respect to ∞ as in §1.2;
$\tau \in \mathrm{Gal}(K/F)$ be the non-trivial element of the Galois group of K/F:
E/F be an elliptic curve with conductor I;
$\mathcal{P}$ be the set of prime numbers associated to E, F, K as in §3.1;
$l \in \mathcal{P}$ be a prime number in $\mathcal{P}$.

As in paragraph (1.12.2), for every prime number $l \neq p$, an isomorphism is fixed between $\mathrm{Gal}(F(E_{l^\infty})/F)$ and a subgroup of $\mathrm{GL}_2(\mathbb{Z}_l)$ by fixing a basis of the corresponding Tate module.

(3.2.2) For each integer $n \geq 1$ and the chosen $l \in \mathcal{P}$, let $\tau_\infty \in \mathrm{Gal}(K(E_{l^n})/F)$ be the unique element of this Galois group satisfying the two conditions:

(a) $\tau_\infty|_{F(E_{l^n})} = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$ that is to say the restriction of τ_∞ to the field extension $F(E_{l^n})/F$ is $\begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$;

(b) $\tau_\infty|_K = \tau$ is the non-trivial element of $\mathrm{Gal}(K/F)$.

The elements τ and τ_∞ have exact order 2.

(3.2.3) For any $\mathbb{Z}[\mathrm{Gal}(K/F)]$-module M on which multiplication by 2 is an isomorphism, we have a decomposition of M as a sum of eigenspaces under the action of τ, the non-trivial element of $\mathrm{Gal}(K/F)$,

$$M \cong M^+ \oplus M^-$$

where M^+ is the submodule of M on which τ acts as 1 and where M^- is similarly the submodule of M on which τ acts as -1.

3.2.4. Definition. (i) For a prime divisor $z \in \Sigma_F$, unramified in the field extension $K(E_{l^n})/F$, let

$$\mathrm{Frob}(z)$$

denote the conjugacy class of $\mathrm{Gal}(K(E_{l^n})/F)$ containing the Frobenius substitutions of the prime divisors above z.

(ii) For $l \in \mathcal{P}$, let $\Lambda^1(n)$ be the set of prime divisors z in Σ_F, of support coprime to ∞ and $\mathrm{Supp}(I)$ and the discriminant of K/F, which satisfy

$$\mathrm{Frob}(z) = [\tau_\infty]$$

where $[\tau_\infty]$ denotes the conjugacy class in $\mathrm{Gal}(K(E_{l^n})/F)$ of τ_∞.
(iii) For $r \geq 0$, let $\Lambda^r(n)$ be the set of effective divisors $z_1 + \ldots + z_r$ on the affine curve Spec A which have r prime components z_i all of which have multiplicity 1 and belong to $\Lambda^1(n)$.

We conventionally put for all $n \geq 1$

$$\Lambda^0(n) = \{0\}$$

which is the set consisting of the zero divisor on C/k.
(iv) Put

$$\Lambda^r = \Lambda^r(1)$$

$$\Lambda(n) = \bigcup_{r \geq 0} \Lambda^r(n)$$

$$\Lambda = \bigcup_{n \geq 1} \Lambda(n).$$

We have

$$\Lambda^r(n) = \left\{ \begin{array}{lcc} z_1 + \ldots + z_r \in \mathrm{Div}_+(A) & | & z_1, \ldots, z_r \text{ are distinct prime divisors such} \\ & | & \text{that for all } i,\ z_i \text{ is prime to } \infty,\ \mathrm{Supp}(I) \\ & | & \text{and the discriminant of } K/F \text{ and} \\ & | & \mathrm{Frob}(z_i) = [\tau_\infty] \text{ on } K(E_{l^n}) \end{array} \right\}.$$

The set Λ^r has a decreasing filtration

$$\Lambda^r = \Lambda^r(1) \supseteq \Lambda^r(2) \supseteq \Lambda^r(3) \supseteq \ldots.$$

3.2.5. Remarks. (1) The prime divisors in $\Lambda(n)$ are infinite in number, by the Chebotarev density theorem, remain prime in the field extension K/F, and their liftings to K split completely in $K(E_{l^n})/K$. Furthermore, E has good reduction at all prime divisors of $\Lambda(n)$.

Note that the prime number $l \in \mathcal{P}$ is considered to be fixed and the sets $\Lambda^r(n)$ depend on l.

(2) The set $\Lambda(n)$ is defined for any prime number l in $\mathcal{P}$ and contains only effective divisors on Spec A consisting of sums of distinct prime divisors whose corresponding Frobenius conjugacy classes in $\mathrm{Gal}(K(E_{l^n})/F)$ are all the same and equal to $[\tau_\infty]$. This unique Frobenius conjugacy class is of a special kind, in particular its elements have order 2.

(3.2.6) For any prime number λ distinct from the characteristic of F, let

$$\rho : \mathrm{Gal}(F^{\mathrm{sep}}/F) \to \mathrm{Aut}_{\mathbb{Q}_\lambda}(T_\lambda(E) \otimes_{\mathbb{Z}_\lambda} \mathbb{Q}_\lambda)$$

denote the galois representation on the λ-adic Tate module $T_\lambda(E)$ of the elliptic curve E; that is to say

$$T_\lambda(E) \otimes_{\mathbb{Z}_\lambda} \mathbb{Q}_\lambda = H^1_{\text{ét}}(E \otimes_F F^{\mathrm{sep}}, \mathbb{Q}_\lambda)^*$$

where $*$ denotes the dual $\mathbb{Q}_\lambda$-vector space.

If z is a prime of F let I_z be an inertia subgroup of $\mathrm{Gal}(F^{\mathrm{sep}}/F)$ at z; let

$$a_z = \mathrm{Tr}(\rho(\mathrm{Frob}(z)) \mid (T_\lambda(E) \otimes_{\mathbb{Z}_\lambda} \mathbb{Q}_\lambda)^{I_z}).$$

That is to say, a_z is the trace of the Frobenius at z on the part of the Tate module invariant under I_z. Then we have $a_z \in \mathbb{Z}$ (see [1, Chapter 5, Examples 5.3.18(1)]).

3.2.7. Lemma. *Suppose that $z \in \Sigma_F$ is a prime divisor of F and $l \in \mathcal{P}$; write $\kappa(z)$ for the residue field at z. .*
(i) *If $z \in \Lambda^1(n)$ then we have $a_z \equiv |\kappa(z)| + 1 \equiv 0 \pmod{l^n}$.*
(ii) *If $z \in \Lambda^1(n)$ and $\mathcal{E}_{0,z}$ denotes the closed fibre over z of the Néron model of E/F and y is the prime divisor of K lying over z then we have group isomorphisms, for $\delta = +1$ or -1,*

$$\mathcal{E}_{0,z}(\kappa(y))^\delta_{l^n} \cong \mathcal{E}_{0,z}(\kappa(z))_{l^n} \cong \mathbb{Z}/l^n\mathbb{Z}.$$

Proof. (i) As a_z is the trace of a Frobenius above z on the Tate module of E/F, we have by the Grothendieck-Lefschetz trace formula, where $\kappa(z)$ is the residue field of F at z and $\mathcal{E}_{0,z}$ is the closed fibre over z of the Néron model of E/F,

$$a_z = |\kappa(z)| + 1 - |\mathcal{E}_{0,z}(\kappa(z))|.$$

On the one hand, let K' be the field $K(E_{l^n})$. The characteristic polynomial of the Frobenius $\mathrm{Frob}_{K'/F}(z)$ above z acting on the l-adic Tate module of E/F is equal to

$$X^2 - a_z X + |\kappa(z)|$$

On the other hand as $z \in \Lambda^1(n)$, the prime z is a place of good reduction of E/F (see remark 3.2.5(1)). Therefore the characteristic polynomial of the Frobenius $\mathrm{Frob}_{K'/F}(z) = [\tau_\infty]$ above z acting on E_{l^n}, the group scheme of l^n-torsion points of E, is equal to by (3.2.2)(a)

$$X^2 - 1 \pmod{l^n}.$$

Comparing these two quadratic polynomials modulo l^n proves the congruences in part (i) of the lemma.

(ii) Let y be the unique place of K lying over the place z of F where $z \in \Lambda^1(n)$. Then $\kappa(y)$ is a quadratic extension of $\kappa(z)$. Furthermore, as $\mathrm{Frob}_{K'/F}(z) = [\tau_\infty]$ where τ_∞ has order 2 (see (3.2.2), remark 3.2.5(2)), the prime y splits completely in the extension K'/K. The map of reduction modulo a prime of K' over z

$$E(K')_{l^n} \to \mathcal{E}_{0,z}(\kappa(y))_{l^n}$$

is an isomorphism. Hence we have

$$\mathcal{E}_{0,z}(\kappa(y))_{l^n} \cong \left(\frac{\mathbb{Z}}{l^n\mathbb{Z}}\right)^2.$$

As l is an odd prime number and the roots ± 1 of the characteristic polynomial $X^2 - 1$ of τ_∞ on E_{l^n} are rational over the prime field $\mathbb{Z}/l\mathbb{Z}$, the action of τ_∞ on $\mathcal{E}_{0,z}(\kappa(y))_{l^n}$ decomposes into a sum over the eigenspaces of τ_∞. Hence we have for $\delta = \pm 1$

$$\mathcal{E}_{0,z}(\kappa(y))^\delta_{l^n} \cong \mathbb{Z}/l^n\mathbb{Z}.$$

Furthermore, we have

$$\mathcal{E}_{0,z}(\kappa(y))^+ = \mathcal{E}_{0,z}(\kappa(z)).$$

The result follows from this. □

3.2.8. Remarks. (1) Let $z \in \Sigma_F$ be a prime divisor which is coprime to ∞, $\mathrm{Supp}(I)$ and the discriminant of K/F and which is inert in the field extension K/F. Then it can be shown, in a similar way to the proof of Lemma 3.2.7, that $z \in \Lambda^1(n)$ if and only if

$$a_z \equiv |\kappa(z)| + 1 \equiv 0 \pmod{l^n}$$

and the Frobenius $\mathrm{Frob}(z)$ does *not* act as a homothety on $T_l(E) \otimes_{\mathbb{Z}_l} \mathbb{Z}/l^n\mathbb{Z}$ where a_z is the trace of the Frobenius $\mathrm{Frob}(z)$ on the Tate module as in §3.2.6.

(2) Fix a prime $l \in \mathcal{P}$. For two integers $a, b \in \mathbb{Z}$ denote by $v_l(a, b)$ the valuation at l of the greatest common divisor of a and b. For a prime divisor z of Spec A define $\alpha(z)$ by the equation

$$\alpha(z) = v_l(|\kappa(z)| + 1, a_z)$$

where as in §3.2.6

$$a_z = |\kappa(z)| + 1 - |\mathcal{E}_0(\kappa(z))|$$

and where $\mathcal{E}_0$ is the closed fibre of the Néron model of E at z.

The prime divisors z of Spec A in the set Λ^1 have the property that they are coprime to I, remain prime in K/F, and satisfy

$$\alpha(z) = v_l(|\kappa(z)| + 1, a_z) \geq 1.$$

(3) If $r > 0$ and $c \in \Lambda^r$ we write

$$\alpha(c) = \min_{z \in \mathrm{Supp}(c)} \alpha(z), \quad \alpha(0) = +\infty.$$

We have the following equation

$$\Lambda^r(n) = \{c \in \Lambda^r \mid \alpha(c) \geq n\}$$

where n is any integer ≥ 1; $\alpha(c)$ is the greatest integer n such that $c \in \Lambda^r(n)$.

The set Λ^r is then equipped with a decreasing filtration

$$\Lambda^r = \Lambda^r(1) \supseteq \Lambda^r(2) \supseteq \Lambda^r(3) \supseteq \dots$$

(4) In the notation of the monograph [1, Chapter 7, §7.11], we have $\Lambda(n) = \mathcal{D}_{l^n}$

3.3. A refined Hasse principle for finite group schemes

The finite group schemes in question are E_{l^n}/F and the Hasse principle concerns the localization at places of F of finite groups of their principal homogenous spaces.

(3.3.1) Suppose that

E is an elliptic curve defined over F;
K is an imaginary quadratic field over F with respect to ∞:
$l \in \mathcal{P}$ is a, necessarily odd, prime number in the set $\mathcal{P}$ (see §3.1);
$n \geq 1$ is an integer;
L_n is the field $K(E_{l^n})$ which is galois over F.

3.3.2 Lemma. *The restriction map from K to L_n induces an isomorphism*

$$H^1(K, E_{l^n}(L_n)) \xrightarrow{\cong} \mathrm{Hom}_{\mathrm{Gal}(L_n/K)}(\mathrm{Gal}(L_n^{\mathrm{ab}}/L_n), E_{l^n}(L_n))$$

where L_n^{ab} is the maximal separable abelian extension of L_n.

Proof. The Hochschild-Serre spectral sequence, where as in (3.3.1) $L_n = K(E_{l^n})$,

$$H^p(L_n/K, H^q(L_n, E_{l^n}(L_n))) \Rightarrow H^{p+q}(K, E_{l^n}(L_n))$$

gives rise to a short exact sequence of low degree terms

$$0 \to H^1(L_n/K, E_{l^n}(L_n)) \to H^1(K, E_{l^n}(L_n)) \to H^1(L_n, E_{l^n}(L_n))^{\mathrm{Gal}(L_n/K)} \to$$

$$\to H^2(L_n/K, E_{l^n}(L_n)) \to H^2(K, E_{l^n}(L_n)).$$

The two cohomology groups $H^1(L_n/K, E_{l^n}(L_n))$ and $H^2(L_n/K, E_{l^n}(L_n))$ are zero by the definition of $\mathcal{P}$ (Definition 3.1.3(b)); hence this exact sequence shows that restriction map of the lemma is an isomorphism.

$$H^1(K, E_{l^n}(L_n)) \to H^1(L_n, E_{l^n}(L_n))^{\mathrm{Gal}(L_n/K)}$$

is an isomorphism of $\mathrm{Gal}(L_n/F)$-modules. The isomorphism of the lemma follows immediately. □

3.3.3. Proposition. *If S is a finite subgroup of $H^1(K, E_{l^n})$ then there is a finite abelian extension $L_{S,n}$ of L_n and an isomorphism of $\mathrm{Gal}(L_n/K)$-modules*

(3.3.4)
$$\mathrm{Gal}(L_{S,n}/L_n) \cong \mathrm{Hom}(S, E_{l^n}(L_n))$$

$$\sigma \mapsto \phi_\sigma$$

and an isomorphism of abelian groups (if S is a $\mathbb{Z}[\mathrm{Gal}(K/F)]$-module then an isomorphism of $\mathbb{Z}[\mathrm{Gal}(K/F)]$-modules)

$$S \cong \mathrm{Hom}_{\mathrm{Gal}(L_n/K)}(\mathrm{Gal}(L_{S,n}/L_n), E_{l^n}(L_n)).$$

For the proof of Proposition 3.3.3, see [1, Chapter 7, Corollary 7.18.10]. Note that the set of prime numbers $\mathcal{P} \setminus F$ of [1, Chapter 7, (7.18.1) and corollary 7.18.10] coincides with the set $\mathcal{P}$ of prime numbers defined in §3.1 above. □

(3.3.5) For a prime divisor v of K, denote by

$$\mathrm{res}_v : H^1(K, E_{l^n}) \longrightarrow H^1(K_v, E_{l^n})$$

the restriction homomorphism at v where K_v is the completion of K at v.

Let S be a finite subgroup of $H^1(K, E_{l^n})$. For $s \in S$ and v a place of K, we have in the notation of Proposition 3.3.3 applied to the subgroup S

$$\mathrm{res}_v(s) = 0 \Leftrightarrow \phi_\sigma(s) = 0 \text{ for all } \sigma \in D_{v'}$$

where v' is a prime divisor of L_n above v and $D_{v'}$ is the decomposition group of a prime divisor of $L_{S,n}$ above v' and where $\phi_\sigma \in \mathrm{Hom}(S, E_{l^n}(L_n))$ as in (3.3.4).

(3.3.6) Let $\tau_\infty \in \mathrm{Gal}(L_n/F)$ be the element of order 2 of paragraph (3.2.2). The element τ_∞ acts as -1 on the l^nth roots of unity and preserves the Weil pairing on E_{l^n} (referring to the exact sequence (1.11.3), $\det(\tau_\infty) = -1$ acts on the l^nth roots of unity).

We obtain (see paragraph (3.2.3)) as l is odd the decompositions into eigenspaces under the action of τ_∞

$$H^1(K, E_{l^n}) \cong H^1(K, E_{l^n})^+ \oplus H^1(K, E_{l^n})^-$$

$$E_{l^n}(L_n) \cong E_{l^n}(L_n)^+ \oplus E_{l^n}(L_n)^-.$$

We obtain that the $+1$ eigenspace

$$\mathrm{Hom}(H^1(K, E_{l^n}), E_{l^n}(L_n))^+$$

is isomorphic to the profinite group of $\mathbb{Z}[\tau_\infty]$-homomorphims from $H^1(K, E_{l^n})$ to $E_{l^n}(L_n)$ and hence we obtain an isomorphism between this τ_∞-invariant subgroup and the profinite Pontrjagin dual of the discrete torsion abelian group $H^1(K, E_{l^n})$ namely we have an isomorphism, where a basis of $T_l(E)$ is fixed as in (1.12.2),

$$\mathrm{Hom}(H^1(K, E_{l^n}), E_{l^n}(L_n))^+ \ \cong \ H^1(K, E_{l^n})^*$$

where the Pontrjagin dual is given by

$$H^1(K, E_{l^n})^* = \mathrm{Hom}(H^1(K, E_{l^n}), \mathbb{Q}/\mathbb{Z}).$$

For a finite subgroup S of $H^1(K, E_{l^n})$ there is similarly an isomorphism

$$\widehat{S} \cong \mathrm{Hom}(S, E_{l^n}(L_n))^+.$$

3.3.7. Proposition. *Let S be a finite subgroup of $H^1(K, E_{l^n})$ and let $\chi \in \widehat{S}$. For any integer $t \geq n$, there is a set of positive Dirichlet density of prime divisors $z \in \Lambda^1(t)$ of F unramified in $L_t = K(E_{l^t})$ such that*

$$\chi = \phi_{\mathrm{Frob}(z^\times)}$$

for some prime divisor $z^\times$ of L_t lying above z, where $\phi_{\mathrm{Frob}(z^\times)} \in \mathrm{Hom}(S, E_{l^t}(L_t))^+$ is as in (3.3.4).

Proof. By Lemma 3.1.5, there is an injection of cohomology groups for all $t \geq n$

$$H^1(K, E_{l^n}) \to H^1(K, E_{l^t})$$

obtained from the inclusion of finite group schemes $E_{l^n} \subseteq E_{l^t}$. The finite subgroup S is then a subgroup of $H^1(K, E_{l^t})$ for all $t \geq n$. Applying Proposition 3.3.3 to S as a subgroup of $H^1(K, E_{l^t})$ we obtain the abelian extension $L_{S,t}/L_t$ where we write $L_t = K(E_{l^t})$.

By Proposition 3.3.3 there is an element $\sigma \in \mathrm{Gal}(L_{S,t}/L_t)$ such that

$$\chi = \phi_\sigma.$$

We have

$$\mathrm{Gal}(L_{S,t}/L_t)^+ \cong \widehat{S}$$

from the isomorphism of (3.3.4) and the isomorphism of paragraph (3.3.6)

$$\widehat{S} \cong \mathrm{Hom}(S, E_{l^t}(L_t))^+.$$

In particular we have $\phi_\sigma^{\tau_\infty} = \phi_\sigma$. As the order of $\mathrm{Gal}(L_{S,t}/L_t)$ is odd, we then have $\sigma = \rho^{\tau_\infty}.\rho$ for some $\rho \in \mathrm{Gal}(L_{S,t}/L_t)$. By the Chebotarev density theorem there is a set of positive Dirichlet density of prime divisors $z \in \Sigma_F$ of F such that $\mathrm{Frob}(z)$ in $\mathrm{Gal}(L_{S,t}/F)$ contains $\tau_\infty\rho$ and where z unramified in $L_{S,t}/F$. Note the the finitely many prime divisors ramified in the field extension $L_{S,t}/F$ depend only on S and not on t.

Since the restriction of $\tau_\infty\rho$ to L_t is τ_∞, we have $z \in \Lambda^1(t)$. We then have that z has residue class extension degree 2 in L_t/F for any prime divisor above z in L_t. Hence for any $z^\times$, a prime divisor of L_t lying above z, we have

$$\mathrm{Frob}(z^\times) = (\tau_\infty\rho)^2 = \rho^{\tau_\infty}.\rho = \sigma. \quad \square$$

3.3.8. Proposition. *Let S be a finite subgroup of $H^1(K, E_{l^n})$ and let $\chi \in \widehat{S}$. Then for any integer $t \geq 1$ there is a set of positive Dirichlet density of prime divisors $z \in \Lambda^1(t)$ such that for the prime divisor y of K lying over z, there is a commutative diagram of group homomorphisms*

$$\begin{array}{cc} S & \xrightarrow{\mathrm{res}_y} \quad \mathrm{res}_y(S) \\ \chi \searrow & \cong\downarrow \psi \\ & \chi(S) \end{array}$$

where ψ is an isomorphism of finite cyclic groups.

Proof. As $\Lambda^1(m)$, $m \geq 1$, is a decreasing filtration of $\Lambda^1(1)$, we may assume that $t \geq n$. By Lemma 3.1.5, there is an injection of cohomology groups for all $t \geq n$

$$H^1(K, E_{l^n}) \to H^1(K, E_{l^t})$$

obtained from the inclusion of finite group schemes $E_{l^n} \subseteq E_{l^t}$. The finite subgroup S is then a subgroup of $H^1(K, E_{l^t})$ for all $t \geq n$.

Select the prime divisor $z \in \Sigma_F$ as in Proposition 3.3.7 applied to S, χ and where the divisors z are different from the finitely many prime divisors of F where the cohomology classes of the finite group S ramify, that is to say the finitely many prime divisors where the field extension $L_{S,t}/F$ of (3.3.4) is ramified. For such a z, let y be the prime divisor of K above z.

The decomposition group of y in $\mathrm{Gal}(L_{S,t}/K)$ is generated by the Frobenius element $\mathrm{Frob}(y)$ as the field extension $L_{S,t}/K$ is unramified at y. By paragraph (3.3.5) we then have

$$\ker(S \xrightarrow{\mathrm{res}_y} H^1(K_y, E_{l^t})) = \ker(\chi).$$

That is to say, the kernel in S of the restriction homomorphism at y of the classes of S is equal to $\ker(\chi)$. Hence the two finite cyclic groups $\mathrm{res}_y(S)$, $\chi(S)$ are isomorphic and we obtain the isomorphism ψ of the commutative diagram of the corollary. □

3.3.9. Corollary. *We have*

$$\begin{array}{ll} \Lambda^r(n) & \textit{is infinite for all integers } \ r, n \geq 1; \\ \Lambda^0(n) = \{0\} & \textit{for all } n \geq 1. \end{array}$$

Proof. We have from Proposition 3.3.8 that $\Lambda^1(n)$ has infinitely many elements for all integers $n \geq 1$. As $\Lambda^r(n)$ consists of all sums of r distinct prime divisors of $\Lambda^1(n)$ (see Definition 3.2.4) and as $\Lambda^0(n) = \{0\}$, the corollary then follows. □

3.3.10. Remark. Proposition 3.3.8 implies that, under the hypotheses of the proposition, the kernel of the natural homomorphism

$$\psi : H^1(K, E_{l^n}) \to \prod_{z \in \Lambda^1(n)} H^1(K_{y(z)}, E_{l^n})$$

is zero, where $y(z)$ is the place of K over $z \in \Lambda^1(n)$. To show this, it is sufficient to apply the proposition to a faithful character χ of the finite cyclic group S generated by any element of $H^1(K, E_{l^n})$.

The *Hasse principle* for the group scheme E_{l^n} and the set of prime divisors $\Lambda^1(n)$ is precisely that the kernel of the homomorphism ψ is zero. It says that a prinicpal homogeneous space of E_{l^n} which is locally trivial at all places of F in $\Lambda^1(n)$ must be globally trival.

In this way, the proposition is a refined form of the Hasse principle for the finite group scheme E_{l^n}. It would be interesting to know to what other group schemes the localization properties of finite subgroups of cohomology groups given in Proposition 3.3.8 also hold.

[For more details on the Hasse principle, see [9, pp.142-150].]

3.4. Drinfeld-Heegner points and the cohomology classes $\gamma_n(c), \delta_n(c)$

(3.4.1) Let

- E/F be an elliptic curve equipped with an origin, that is to say E/F is a 1-dimensional abelian variety;
- I be the ideal of A which is the conductor of E/F without the component at ∞;
- $\epsilon = \pm 1$ be the sign in the functional equation of the L-function of the elliptic curve E/F.

Assume that (see also [1, Chapter 4, §§4.3, 4.7, or Chapter 7, §7.6]):

- (α) E/F has split Tate multiplicative reduction at ∞ (see §1.7 or [1, Chapter 4, §4.7]);
- (β) K is an (separable) imaginary quadratic field extension of F, with respect to ∞, such that all primes dividing the conductor I split completely in K.

The hypothesis (α) implies that E/F is covered by the Drinfeld modular curve $X_0^{\mathrm{Drin}}(I)$ (see (3.4.4) below). The hypothesis (β) ensures the existence of Drinfeld-Heegner points on $X_0^{\mathrm{Drin}}(I)$. The two hypotheses together ensure that there are Drinfeld-Heegner points on the elliptic curve E/F. Note that from (3.4.7) to the

end of this section §3.4, it is assumed that $K \neq F \otimes_{\mathbb{F}_q} \mathbb{F}_{q^2}$ where $\mathbb{F}_q$ is the exact field of constants of F.

In this section we detail this construction of Drinfeld-Heegner points as well as defining the cohomology classes $\gamma_n(c), \delta_n(c)$.

(3.4.2) Let λ be any prime number of $\mathbb{Z}$ distinct from the characteristic of F. Let ρ be the 2-dimensional λ-adic representation of $\mathrm{Gal}(F^{\mathrm{sep}}/F)$ corresponding to E where F^{sep} is the separable closure of F; that is to say ρ is the continuous homomorphism

$$\rho : \mathrm{Gal}(F^{\mathrm{sep}}/F) \to \mathrm{End}_{\mathbb{Q}_\lambda}(H^1_{\text{ét}}(E \otimes_F F^{\mathrm{sep}}, \mathbb{Q}_\lambda)).$$

For each place v of F put

$$a_v = \mathrm{Tr}(\rho(\mathrm{Frob}(v))|\ H^1_{\text{ét}}(E \otimes_F F^{\mathrm{sep}}, \mathbb{Q}_\lambda)^{T_v})$$

where T_v is the inertia subgroup of $\mathrm{Gal}(F^{\mathrm{sep}}/F)$ over v. The representation ρ satisfies

$$a_v \in \mathbb{Z} \text{ for all } v \in \Sigma_F$$

(see [1, Chapter 5, example 5.3.18]).

(3.4.3) Let

B be the integral closure of A in K;
O_c be the order of K, with respect to A, and of conductor c for any divisor $c \in \mathrm{Div}_+(A)$, (see [1, Chapter 2, §2.2]);
τ be the non-trivial element of $\mathrm{Gal}(K/F)$;
$IB = I_1 I_2$ be a factorisation of ideals of B where I_1, I_2 are ideals of B such that $I_2^\tau = I_1$ where I is the ideal of A which is the conductor of E/F without the place at ∞, as in (3.4.1); such a factorisation exists because of the hypothesis that the prime ideal components of I split completely in K/F (by (3.4.1)(β)).

As in §1.4, $K[c]$ denotes the ring class field of K of conductor $c \in \mathrm{Div}_+(A)$. In particular $K[0]$ is the Hilbert class field of K, that is to say $K[0]$ is the maximal unramified abelian extension of K which is split completely at ∞.

(3.4.4) Let $\mathcal{H}(\rho)$ be the Heegner module of ρ and K/F with exceptional set of primes those dividing I and the place ∞ with coefficients in $\mathbb{Z}$ (see [1, Chapter 5, §5.3]), where I is the conductor of E, without the component at ∞.

As in §1.7 (see also [1, Chapter 4, §4.7] and [1, Appendix B]), there is a finite surjective morphism of curves over F under the hypothesis (α)

$$\pi : X_0^{\mathrm{Drin}}(I) \to E.$$

We may translate π in the group scheme E so that $\pi^{-1}(0)$ consists of at least one cusp of $X_0^{\mathrm{Drin}}(I)$ (as in [1, Chapter 4, (4.8.1)]); this rigidifies the map π. The cusps of the modular curve $X_0^{\mathrm{Drin}}(I)$ generate a torsion subgroup of the jacobian of this curve [1, Chapter 2, Theorem 2.4.9].

Let a be a divisor class in the Picard group $\mathrm{Pic}(O_c)$ of the order O_c. Assume that c and I are coprime. Then there is a Drinfeld-Heegner point

$$(a, I_1, c) \in X_0^{\mathrm{Drin}}(I)(K[c])$$

which is a non-cuspidal point of $X_0^{\mathrm{Drin}}(I)$ and is rational over the ring class field $K[c]$.

This point (a, I_1, c) is constructed as follows. Fix an embedding $K \to \widehat{\overline{F}}_\infty$ where $\widehat{\overline{F}}_\infty$ is the completion of the algebraic closure of F_∞ which is the completion of F at ∞. Let L be a projective O_c-module of rank 1 in the class a and contained as a lattice in $\widehat{\overline{F}}_\infty$. Then $I_1(O_c) = I_1 \cap O_c$ is an invertible ideal of O_c and $L' = I_1(O_c)^{-1}L$ is a projective O_c module of rank 1 contained as a lattice in $\widehat{\overline{F}}_\infty$. Let D and D' be the rank 2 Drinfeld modules for A over the field $\widehat{\overline{F}}_\infty$ corresponding respectively to the lattices L and L'. Then D and D' have general characteristic and complex multiplication by O_c. The inclusion of O_c-modules $L \subset L'$ corresponds to an I-cyclic isogeny $f : D \to D'$, as its kernel is isomorphic as an A-module to $O_c/I_1(O_c) \cong A/I$. The pair $(D, \ker(f))$ defines the point (a, I_1, c) on $X_0^{\mathrm{Drin}}(I)(\widehat{\overline{F}}_\infty)$. That this point (a, I_1, c) is defined over $K[c]$ results from the main theorem of complex multiplication (see [1, Chapter 4, §4.3] for more details).

(3.4.5) The image

$$\pi(a, I_1, c) \in E(K[c])$$

is a Drinfeld-Heegner point of E rational over the ring class field $K[c]$ and is written in the notation of [1, Chap. 4, §4.8]

$$(a, I_1, c, \pi) \in E(K[c]).$$

By [1, Chap. 5, example 5.3.18] there is a homomorphism of discrete $\mathrm{Gal}(K^{\mathrm{sep}}/K)$-modules

$$\begin{aligned} \mathcal{H}(\pi): \quad \mathcal{H}(\rho)^{(0)} &\to E(F^{\mathrm{sep}}) \\ < a, c > &\mapsto (a, I_1, c, \pi) \end{aligned}$$

for all $c \in \mathrm{Div}_+(A)$ coprime to I, $a \in \mathrm{Pic}(O_c)$. The image of this homomorphism $\mathcal{H}(\pi)$ consists of the $\mathbb{Z}$-linear combinations of Drinfeld-Heegner points of E rational over all the ring class fields $K[c]$ for all c.

Let $< 0, 0 >$ be the element of the Heegner module $\mathcal{H}(\rho)^{(0)}$ given by the principal class of $\mathrm{Pic}(B)$, where B is the integral closure of A in K. Let

$$(0, I_1, 0, \pi) = \mathcal{H}(\pi)(< 0, 0 >)$$

be the corresponding Drinfeld-Heegner point of $E(K[0])$ (see [1, Chapter 4, (4.8.2)]). Let

$$P_0 = \mathrm{Tr}_{K[0]/K}(0, I_1, 0, \pi) \in E(K). \tag{3.4.6}$$

That is to say, P_0 is the trace from $K[0]$ to K of the point $(0, I_1, 0, \pi)$; the point P_0 belongs to $E(K)$ and the point $(0, I_1, 0, \pi)$ belongs to $E(K[0])$.

(3.4.7) We now impose for the rest of this section the hypothesis that $K \neq F \otimes_{\mathbb{F}_q} \mathbb{F}_{q^2}$ where $\mathbb{F}_q$ is the exact field of constants of F, that is to say K is not obtained from F by ground field extension.

Let $c \in \Lambda(1)$ that is to say c is a sum of distinct prime divisors of $\Lambda^1(1)$ with multiplicity 1. Let

$$y_c = \pi(0, I_1, c) \in E(K[c]).$$

so that $y_c = (0, I_1, c, \pi)$. The field $K[0]$ is the Hilbert class field of K. Let

$$G_c = \mathrm{Gal}(K[c]/K[0]).$$

As $K \neq F \otimes_{\mathbb{F}_q} \mathbb{F}_{q^2}$ by hypothesis, we have that $B^* = A^*$, where A^*, B^* are the unit groups of A, B, and hence (by [1, Chap. 2, (2.3.8), p.19]) there is a group isomorphism

$$G_c \cong \prod_z G_z$$

where the product runs over the prime divisors z in the support of c and

$$G_z = \mathrm{Gal}(K[c]/K[c - z])$$

is a cyclic group of order $|\kappa(z)|+1$ as z is inert in K/F (by [1, Chapter 2, (2.3.12), p.20]). Fix a generator σ_z of the cyclic group G_z for all prime divisors $z \in \Lambda^1(1)$.

(3.4.8) Write for any prime divisor $z \in \Lambda^1(1)$ of F, where $D_z \in \mathbb{Z}[G_z]$,

$$D_z = -\sum_{i=1}^{|\kappa(z)|} i.\sigma_z^i.$$

Here D_z is the Kolyvagin element of the map $h_z : G_z \to \mathbb{Z}$ where $\sigma_z^{-i} \mapsto -i$ for all $0 \leq i \leq |\kappa(z)|$. Note the minus sign here which agrees with the Kolyvagin elements of [1, Chap. 5, pp. 175-178].

(3.4.9) For any divisor $c \in \Lambda(1)$, put

$$D_c = \prod_{z \in \ \mathrm{Supp}(c)} D_z$$

where $D_c \in \mathbb{Z}[G_c]$. Then D_c is the Kolyvagin element of the map $h : G_c \to \mathbb{Z}$ given by $h = \prod_{z \in \mathrm{Supp}(z)} h_z$. Let

$$\mathcal{G}_c = \mathrm{Gal}(K[c]/K)$$

where there is an exact sequence of finite abelian groups

$$0 \longrightarrow G_c \longrightarrow \mathcal{G}_c \longrightarrow \mathrm{Gal}(K[0]/K) \longrightarrow 0.$$

Let $\mathcal{S}$ be a set of coset representatives for G_c in $\mathcal{G}_c$. Define the point $P_c \in E(K[c])$ by, where $y_c = \pi(0, I_1, c)$ is the element of $E(K[c])$ in (3.4.7),

$$P_c = \sum_{s \in \mathcal{S}} s D_c y_c.$$

(3.4.10) Suppose now that $c \in \Lambda(n)$. We write $P_c \pmod{l^n}$ for the image of P_c in the quotient group ${}_{l^n}E(K[c])$. Then we have that $P_c \pmod{l^n}$ belongs to

$$P_c \pmod{l^n} \ \in \left({}_{l^n}E(K[c]) \right)^{\mathcal{G}_c}.$$

This inclusion follows immediately from the formula in $\mathbb{Z}[G_c]$

$$(\sigma_z - 1) D_z = -|G(c/c - z)| + \sum_{g \in G(c/c-z)} g$$

for all $z \in \mathrm{Supp}(c)$ and that $|G(c/c-z)| = |\kappa(z)| + 1$ is divisible by l^n for all $z \in \mathrm{Supp}(c)$ (for a detailed proof of this inclusion $P_c \pmod{l^n} \ \in ({}_{l^n}E(K[c]))^{\mathcal{G}_c}$ see [1, Chapter 7, Lemma 7.14.9 and Lemma 7.14.11]). Furthermore, we have where y_0 is defined in (3.4.7) and where this notation P_0 agrees with that of (3.4.6)

$$P_0 = \mathrm{Tr}_{K[0]/K}(y_0) \in E(K).$$

[The point $P_c \pmod{l^n}$ in $E(K[c])/l^n E(K[c])$ coincides with the point denoted P_c in the book [1, Chapter 7, Notation 7.14.10(iii) and Lemma 7.14.11].]

(3.4.11) Let S be the ring $\mathbb{Z}/m\mathbb{Z}$ where m is any non-zero integer. The morphism of multiplication by m on the elliptic curve E then provides for any divisor c of $\mathrm{Div}_+(A)$ prime to I the following commutative diagram with exact rows and an exact right-hand column (see [1, Chap. 7, §7.14.5]):

(3.4.12)

$$
\begin{array}{ccccccccc}
 & & & & & & 0 & & \\
 & & & & & & \downarrow & & \\
 & & & & & & H^1(K[c]/K, E(K[c]))_m & & \\
 & & & & & \overset{j}{} & \inf\downarrow & & \\
0 \to & {}_mE(K) & \to & H^1(K, E_m) & \to & H^1(K, E)_m & \to & 0 & \\
 & \downarrow & & \text{res}\downarrow \text{quasi}-\text{isom.} & & \text{res}\downarrow & & & \\
0 \to & ({}_mE(K[c]))^{\mathcal{G}_c} & \overset{\partial}{\to} & H^1(K[c], E_m)^{\mathcal{G}_c} & \to & H^1(K[c], E)_m^{\mathcal{G}_c} & & & \\
 & f\uparrow & & & & & & & \\
 & (\mathcal{H}_{c,S}^{(0)})^{\mathcal{G}_c} & & & & & & &
\end{array}
$$

Here $(\mathcal{H}_{c,S}^{(0)})$ is the c-component Heegner module with coefficients in the ring S (see [1, Chap. 7, (7.11.3), and Chap. 5, §5.3]).

Let $\mathcal{E}$ be the finite exceptional set of prime numbers of Propositions 1.10.1 and 1.10.2. The middle restriction homomorphism here in (3.4.12) is a quasi-isomorphism of quasi-groups in $[\mathbf{N}^{(p)}]_{\mathbb{Z}}$ where the finite exceptional set of prime numbers is $\mathcal{E}$ and is independent of c (Propositions 1.10.1 and 1.10.2, or alternatively [1, Chap. 7, Proposition 7.14.2]). In particular this middle restriction homomorphism is an isomorphism for all integers m which are powers of prime numbers of $\mathcal{P}$ (Definition 3.1.3) as $\mathcal{P}$ excludes the finitely many prime numbers of $\mathcal{E}$.

The map f is obtained by sending a generator $< a, c >$ of $\mathcal{H}_{c,S}^{(0)}$ to its image $(a, I_1, c, \pi) \in E(K[c])$ as in (3.4.4) and (3.4.5).

This diagram (3.4.12) then provides the Heegner homomorphism, for all integers $m \in \mathbb{N}$ prime to $\mathcal{E}$,

$$(\mathcal{H}_{c,S}^{(0)})^{\mathcal{G}_c} \to H^1(K, E)$$

whose image belongs to $H^1(K[c]/K, E(K[c]))_m$.

(3.4.13) Take $m = l^n$ where $l \in \mathcal{P}$ so that the middle quasi-isomorphism in diagram (3.4.12) is an isomorphism. From the diagram (3.4.12), define $\gamma_n(c)$ and $\delta_n(c)$ by the formulae

$$P_c \pmod{l^n} \in \left({}_{l^n}E(K[c])\right)^{\mathcal{G}_c};$$

$$\gamma_n(c) \text{ is the image of } P_c \pmod{l^n} \text{ in } H^1(K, E_{l^n});$$

$$\delta_n(c) \text{ is the image of } \gamma_n(c) \text{ in } H^1(K, E)_{l^n}.$$

[This is the same as the construction in [1, Chap. 7, §7.14.10], where we write here $\gamma_n(c), \delta_n(c)$ in place of $\gamma(c), \delta(c)$ to indicate their dependence on the integer n.]

3.4.14. Proposition. (i) *The order of* $\gamma_n(c)$ *is equal to the order of* $P_c \pmod{l^n}$ *in* ${}_{l^n}E(K[c])$.
(ii) *The exponent* t *of the order* l^t *of* $\delta_n(c)$ *is the least integer* t *such that*

$$l^t(P_c \pmod{l^n}) \in \frac{l^n E(K[c]) + E(K)}{l^n E([c])}.$$

Proof. Here $P_c \pmod{l^n}$ denotes the image of $P_c \in E(K[c])$ in ${}_{l^n}E(K[c])$. These results on the orders of $\gamma_n(c)$ and $\delta_n(c)$ follow immediately from their definition and the commutative diagram (3.4.12). □

Chapter 4. Structure of the Tate-Shafarevich group and the Selmer group

4.1. Statement of the main theorems

This section contains no proofs. The main Theorems 4.1.9, 4.1.13, 4.1.14 and 4.1.15 of this paper are finally proved in §§5.3-5.5.

(4.1.1) Throughout this chapter 4, we assume that

E/F is an elliptic curve where F is a global field of characteristic $p > 0$;
I is the ideal of A which is the conductor of E/F without the component at ∞;
K is an imaginary quadratic extension field of F with respect to ∞;
$l \in \mathcal{P}$ is a, necessarily odd, prime number in the set of prime numbers $\mathcal{P}$ (see §3.1);
$\epsilon = \pm 1$ is the sign of the functional equation of the L-function $L(E/F, s)$ of E/F;
τ is the element of order 2 of the galois group $\mathrm{Gal}(K/F)$;
$P_c \in E(K[c])$ are the points defined in (3.4.9) over the ring class fields $K[c]$ for all divisors $c \in \Lambda(1)$, and where P_0 belongs to the group $E(K)$ of K-rational points;
$\alpha(c)$, for a divisor $c \in \Lambda(1)$, is the largest integer n such that $c \in \Lambda(n)$ if $c \neq 0$ and such that $\alpha(0) = +\infty$ (see remarks 3.2.8).

Assume that E, K, F satisfy the following hypotheses (as in [1, Chapter 7, §7.6.1]):

(a) ∞ is a place of F with residue field equal to k;
(b) E/F has split Tate multiplicative reduction at ∞ (see §1.7 or [1, Chapter 4, §4.7]);
(c) K is an (separable) imaginary quadratic field extension of F, with respect to ∞, such that all primes dividing the conductor I split completely in K and $K \neq F \otimes_{\mathbb{F}_q} \mathbb{F}_{q^2}$.

These hypotheses (a),(b),(c) are assumed for the rest of this paper.

(4.1.2) If M is a $\mathrm{Gal}(K/F)$-module denote by M^+ the submodule of M on which τ acts by $+1$ and M^- is similarly the submodule on which τ acts by -1.

(4.1.3) Let G be a finite abelian l-group. The *invariants* of G are the integers

$$r_1 \geq r_2 \geq r_3 \geq \dots$$

such that G decomposes into elementary components

$$G \cong \frac{\mathbb{Z}}{l^{r_1}\mathbb{Z}} \oplus \frac{\mathbb{Z}}{l^{r_2}\mathbb{Z}} \oplus \frac{\mathbb{Z}}{l^{r_3}\mathbb{Z}} \oplus \dots$$

The integers $r_1, r_2, r_3 \dots$ are uniquely determined by G; the integers $l^{r_1}, l^{r_2}, l^{r_3}, \dots$ are also sometimes called the invariants of G.

(4.1.4) As the elliptic curve E is defined over F, the l-power torsion subgroup $Ш(E/K)_{l^\infty}$ of the Tate-Shafarevich group $Ш(E/K)$ of the elliptic curve $E \times_F K$ over K decomposes into eigenspaces

$$Ш(E/K)_{l^\infty} \cong Ш(E/K)^+_{l^\infty} \oplus Ш(E/K)^-_{l^\infty}$$

under the action of the element $\tau \in \mathrm{Gal}(K/F)$.

The Cassels pairing on $Ш(E/K)$ is anti-symmetric and respects this decomposition into eigenspaces; furthermore, the Cassels pairing is non-degenerate if this Tate-Shafarevich group $Ш(E/K)$ is finite; therefore if $Ш(E/K)$ is finite, the invariants of the finite abelian l-group $Ш(E/K)_{l^\infty}$ have even multiplicity.

(4.1.5) Under the hypothesis that the Tate-Shafarevich group $Ш(E/K)_{l^\infty}$ be finite, let

$$N_1 \geq N_3 \geq N_5 \dots$$

and

$$N_2 \geq N_4 \geq N_6 \dots$$

be integers such that

$$N_1, N_1, N_3, N_3, N_5, N_5 \dots$$

are the invariants of the finite abelian l-group $Ш(E/K)^\epsilon_{l^\infty}$, which is the ϵ-eigenspace, and

$$N_2, N_2, N_4, N_4, N_6, N_6 \dots$$

are the invariants of the finite abelian l-group $Ш(E/K)^{-\epsilon}_{l^\infty}$, which is the $-\epsilon$-eigenspace.

That is to say, putting $\nu(r) = (-1)^r\epsilon$ then we have isomorphisms for $r = 1$ or $r = 2$

$$Ш(E/K)^{-\nu(r)}_{l^\infty} \cong \bigoplus_{s\geq 0} (\mathbb{Z}/l^{N_{2s+r}}\mathbb{Z})^2.$$

4.1.6. Definition. Let $m \geq 1$ be an integer and $c \in \Lambda(m)$. Then the point P_c, defined in (3.4.9), belongs to $E(K[c])$ and the point P_0 belongs to $E(K)$.
(1) Write

$$l^m | P_c \quad \text{if} \quad P_c \in l^m E(K[c])$$

and

$$l^m||P_c \text{ if } P_c \in l^m E(K[c]) \setminus l^{m+1}E(K[c]).$$

(2) For any integer $r \geq 0$, let $E(r)$ be the abelian group

$$E(r) = \bigoplus_{c \in \Lambda^r(1)} {}_{A(c)}E(K[c])$$

where the sum runs over all divisors c of $\Lambda^r(1)$ and where $A(c) = l^{\alpha(c)}$. If $r > 0$ the group $E(r)$ is then a direct sum of finite abelian l-groups of the form $E(K[c])/l^{\alpha(c)}E(K[c])$. If $r = 0$ then $E(0)$ is defined conventionally to be $E(K[0])$.

A point $P_c \in E(K[c])$, where $c \in \Lambda^r(1)$, induces an element in ${}_{A(c)}E(K[c])$ and hence an element of $E(r)$ where all its components in $E(r)$ are zero except possibly for that in ${}_{A(c)}E(K[c])$.

(3) Let $P(r)$ be the subgroup of $E(r)$ generated by the images in $E(r)$ of the points P_c for all $c \in \Lambda^r(1)$.

Define M_r for any $r \geq 0$ to be the largest integer $n \geq 0$ such that

$$P(r) \subseteq l^n E(r).$$

If $P(r) = 0$ then there is no such largest integer M_r and we then put

$$M_r = +\infty.$$

4.1.7. Conjecture. For some $r \geq 0$ we have $M_r < +\infty$.

[See [7, Conjecture A], and also [1, Conjecture 7.14.19] for an explanation of this conjecture.]

4.1.8. Remarks. (1) The order of $\gamma_n(c)$ is l^{n-m} for all $n \geq m$ if and only if $l^m||P_c$ by Proposition 3.4.14.

If M_r is finite then there is a divisor $c \in \Lambda^r(1)$ such that $\gamma_{M_r+1}(c) \neq 0$ and $\gamma_{M_r}(c) = 0$; furthermore, $\gamma_{M_r}(c) = 0$ for any $c \in \Lambda^r(1)$ from Proposition 3.4.14(i).

(2) We have $M_0 < +\infty$ if and only if P_0 has infinite order in $E(K)$.

[This equivalence requires that $E(K)$ has no l-torsion; but this requirement holds by the restriction on the prime number l given by Definition 3.1.3(b). See Lemma 5.1.2 below for details.]

(3) We have that $\alpha(c)$ is finite if and only if $c \neq 0$ (see Remarks 3.2.8(2) and (3)). Write if $c \neq 0$

$$\mathrm{ord}_l(P_c) = \begin{cases} \max\{N : l^N | P_c\} & \text{provided that this maximum is } < \alpha(c) \\ +\infty & \text{if } \max\{N : l^N | P_c\} \text{ is } \geq \alpha(c). \end{cases}$$

Write

$$\mathrm{ord}_l(P_0) = \max\{N : l^N | P_0\}$$

which is either an integer or $+\infty$. If $c \neq 0$, we have $\mathrm{ord}_l(P_c) < +\infty$ if and only if $l^{\mathrm{ord}_l(P_c)} || P_c$. and $\mathrm{ord}_l(P_c) < \alpha(c)$.
(4) By definition M_r, for all integers $r \geq 0$, is given by

$$M_r = \min\{\mathrm{ord}_l(P_c) | \ \ c \in \Lambda^r(1)\}.$$

Note that $M_r < +\infty$ if and only if there is $c \in \Lambda^r(1)$ such that

$$\mathrm{ord}_l(P_c) < \alpha(c)$$

with an evident interpretation in the case where $r = 0$, $c = 0$ and $\alpha(0) = +\infty$. Furthermore, $M_r < +\infty$ implies that for this selection of $c \in \Lambda^r(1)$ we would have $l^{M_r} || P_c$ and $c \in \Lambda^r(\alpha(c))$ and $M_r < \alpha(c)$.

Alternatively, if $r > 0$ then we have $M_r < +\infty$ if and only if for some $c \in \Lambda^r(1)$ we have

$$c \in \Lambda^r(m+1) \setminus \Lambda^r(m+2)$$

and

$$\mathrm{ord}_l(P_c) \leq m.$$

4.1.9. Theorem. *Suppose that P_0 has infinite order in $E(K)$, the group of K-rational points of E. Let l be a prime number in $\mathcal{P}$ coprime to the order of* $\mathrm{Pic}(A)$ *the Picard group of A. Then the Tate-Shafarevich group $Ш(E/K)$ is finite and the invariants N_i, with multiplicity* 2*, of the subgroup $Ш(E/K)_{l^\infty}$ are given by*

$$N_i = M_{i-1} - M_i, \quad for \ \ i \geq 1.$$

That is to say, we have the isomorphisms of eigenspaces

$$Ш(E/K)^{\epsilon}_{l^\infty} \cong \prod_{\substack{i \ \mathrm{even} \\ i \geq 0}} (\mathbb{Z}/l^{M_i - M_{i+1}}\mathbb{Z})^2$$

$$Ш(E/K)^{-\epsilon}_{l^\infty} \cong \prod_{\substack{i \ \mathrm{odd} \\ i \geq 0}} (\mathbb{Z}/l^{M_i - M_{i+1}}\mathbb{Z})^2.$$

4.1.10. Remarks. (1) The image of the Drinfeld-Heegner point P_0 in $E(K)/E(K)_{\mathrm{tors}}$ is an eigenvector for τ and its eigenvalue is equal to $-\epsilon$ (by [1, Chapter 4, Theorem 4.8.6, p. 98], see also [1, Chapter 7, Lemma 7.14.11, p. 388]) where $E(K)_{\mathrm{tors}}$ is the torsion subgroup of $E(K)$.
(2) If the Drinfeld-Heegner point P_0 has infinite order in $E(K)$ then by [1, Chapter 7, Theorem 7.7.5 and Remarks 7.7.6(3)] the Tate-Shafarevich group $Ш(E/K)$ is finite and therefore the alternating Cassels pairing on $Ш(E/K)$ is non-degenerate and by §§4.1.4,4.1.5 the invariants of the τ-eigenspaces of $Ш(E/K)_{l^\infty}$ have even multiplicity.

The next corollary is an immediate consequence of Theorem 4.1.9.

4.1.11. Corollary. *Under the hypotheses of Theorem 4.1.9, the integers M_i satisfy*

$$M_i - M_{i+1} \geq M_{i+2} - M_{i+3} \geq 0, \ for\ all\ i \geq 0,$$

and if j is such that

$$M_j = M_{j+1} = M_{j+2}$$

then the sequence $M_j, M_{j+1}, M_{j+2}, M_{j+3} \ldots$ is constant. □

4.1.12. Definition. Let

$$G = G_1 \times \ldots \times G_r$$

be a direct product of finite cyclic groups G_i. The characters $\chi_1, \ldots, \chi_r$ of G form a *triangular basis for the dual* $\widehat{G}$, relative to the product $G = \prod_i G_i$, if they generate $\widehat{G}$ and

$$\chi_i(G_j) = 0, \quad \text{for all } j > i.$$

4.1.13. Theorem. *Assume that P_0 has infinite order in $E(K)$ and $l \in \mathcal{P}$ is a prime number coprime to the order of the Picard group* $\mathrm{Pic}(A)$. *Suppose that the direct product*

$$D = \prod_{i \geq 1} D_i$$

is a maximal isotropic subgroup of $Ш(E/K)_{l^\infty}$ for the Cassels pairing and D_i is a finite cyclic group of order l^{N_i} for all $i \geq 1$ and we have direct products

$$D^\epsilon = \prod_{i \text{ odd}} D_i$$

$$D^{-\epsilon} = \prod_{i \text{ even}} D_i.$$

Then there are effective divisors $c_1 \leq c_2 \leq \ldots$ on F such that $c_i \in \Lambda^i(M_{i-1})$ for all $i \geq 1$ and the characters

$$d \mapsto < d, \delta_{M_{i-1}}(c_i) >_{\text{Cassels}}, \quad for\ all\ i$$

form a triangular basis of characters of D relative to the product $D = \prod_i D_i$.

4.1.14. Theorem. *Assume that P_0 has infinite order in $E(K)$ and $l \in \mathcal{P}$ is coprime to the order of* $\mathrm{Pic}(A)$. *Let*

$$M_\infty = \min_{i \in \mathbf{N}} M_i.$$

Then the group $\mathbb{Z}P_0$ has finite index in $E(K)$ and the highest power of l dividing the index $[E(K) : \mathbf{Z}P_0]$ equals l^{M_0} that is to say

$$|(E(K)/\mathbf{Z}P_0)_{l^\infty}| = l^{M_0}.$$

Furthermore, we have

$$|\text{Ш}(E/K)_{l^\infty}| = l^{2(M_0 - M_\infty)}.$$

4.1.15. Theorem. *Assume that P_0 has infinite order in $E(K)$ and $l \in \mathcal{P}$ is coprime to the order of* $\mathrm{Pic}(A)$. *Then for all integers $m \geq 0$, the natural surjection from the Selmer group to the Tate-Shafarevich group*

$$\pi_m : \mathrm{Sel}_{l^m}(E/K) \longrightarrow \text{Ш}(E/K)_{l^m} \tag{4.1.16}$$

splits and we have isomorphisms of eigenspaces

$$\mathrm{Sel}_{l^m}(E/K)^{\pm} \cong ({}_{l^m}E(K))^{\pm} \oplus \text{Ш}(E/K)^{\pm}_{l^m} \text{ for all } m \geq 0. \tag{4.1.17}$$

Furthermore, we have isomorphisms for all $m \geq 0$

$$\begin{array}{ccc} ({}_{l^m}E(K))^{-\epsilon} & \cong & \mathbb{Z}/l^m\mathbb{Z} \\ ({}_{l^m}E(K))^{\epsilon} & \cong & 0. \end{array} \tag{4.1.18}$$

The next corollary follows immediately from Theorems 4.1.9 and 4.1.15.

4.1.19. Corollary. *Under the hypotheses of Theorem 4.1.15, put*

$$\widehat{N_i} = \min(m, M_{i-1} - M_i)$$

for all i and for all integers m. Then for all integers $m \geq 0$, the invariants of the Selmer $-\epsilon$-eigenspace $\mathrm{Sel}_{l^m}(E/K)^{-\epsilon}$ are $m, \widehat{N_2}, \widehat{N_2}, \widehat{N_4}, \widehat{N_4}, \ldots$ and the invariants of the Selmer ϵ-eigenspace $\mathrm{Sel}_{l^m}(E/K)^{\epsilon}$ are $\widehat{N_1}, \widehat{N_1}, \widehat{N_3}, \widehat{N_3}, \ldots$ □

4.1.20. Remark. The main result on the finiteness of Tate-Shafarevich groups proved in the monograph [1, Chapter 7, Theorem 7.6.5 and Theorem 7.7.5] is required for the proofs of the main results of this paper stated in this section; in particular, this paper does not provide a different proof of finiteness independent of the book [1]. This is also the prinicpal reason why the hypotheses (a), (b), (c) of §4.1.1 are required.

4.2. Cochains for the cohomology classes $\gamma_n(c), \delta_n(c)$

The notation and hypotheses of (4.1.1) and (4.1.2) hold in this section.

4.2.1. Lemma. *Let $c \in \Lambda(n)$.*
(i) *The cohomology class $\gamma_n(c)$ is represented by the cocycle*

$$\sigma \mapsto -\frac{(\sigma-1)P_c}{l^n} + \sigma\frac{P_c}{l^n} - \frac{P_c}{l^n}, \quad \mathrm{Gal}(K^{\mathrm{sep}}/K) \to E(K^{\mathrm{sep}})_{l^n}$$

where $\frac{(\sigma-1)P_c}{l^n}$ is the unique l^n-division point of $(\sigma-1)P_c$ in $E(K[c])$ and P_c/l^n is a fixed l^n-division point of P_c.
(ii) *The cohomology class $\delta_n(c)$ is represented by the cocycle, where $\frac{(\sigma-1)P_c}{l^n}$ is the unique l^n-division point of $(\sigma-1)P_c$ in $E(K[c])$,*

$$\sigma \mapsto -\frac{(\sigma-1)P_c}{l^n}, \quad \mathrm{Gal}(K^{\mathrm{sep}}/K) \to E(K^{\mathrm{sep}}).$$

(iii) *Let $n \geq m$ be positive integers and let $c \in \Lambda(n)$. Then we have*

$$l^m\delta_n(c) = \delta_{n-m}(c)$$

$$l^m\gamma_n(c) = \gamma_{n-m}(c).$$

Proof of Lemma 4.2.1. (i) and (ii) These two formulae for cocycles representing the cohomology classes $\gamma_n(c)$ and $\delta_n(c)$ can be extracted from Step 2 of the proof of [1, Chapter 7, Lemma 7.14.14]. We reprove these formulae here.

From the diagram (3.4.12) the restriction homomorphism, where $\mathcal{G}_c = \mathrm{Gal}(K[c]/K)$,

$$H^1(K, E_{l^n}(K^{\mathrm{sep}})) \to H^1(K[c], E_{l^n}(K^{\mathrm{sep}}))^{\mathcal{G}_c}$$

is an isomorphism as the prime number l belongs to $\mathcal{P}$. The point P_c belongs to $E(K[c])$.

Let $\frac{P_c}{l^n} \in E(K^{\mathrm{sep}})$ be a fixed l^nth division point of P_c, that is to say $\frac{P_c}{l^n}$ is any point which satisfies $l^n(\frac{P_c}{l^n}) = P_c$. Then the cocycle

$$\phi : g \mapsto g(\frac{P_c}{l^n}) - \frac{P_c}{l^n}, \ \mathrm{Gal}(K^{\mathrm{sep}}/K[c]) \to E_{l^n}(K^{\mathrm{sep}}),$$

represents a cohomology class in $H^1(K[c], E_{l^n}(K^{\mathrm{sep}}))^{\mathcal{G}_c}$ which is the image of $P_c \pmod{l^n} \in ({}_{l^n}E(K[c]))^{\mathcal{G}_c}$ under the coboundary map

$$\partial_{l^n} : ({}_{l^n}E(K[c]))^{\mathcal{G}_c} \to H^1(K[c], E_{l^n}(K^{\mathrm{sep}}))^{\mathcal{G}_c}$$

(see the diagram (3.4.12)). The inflation of ϕ to $\mathrm{Gal}(K^{\mathrm{sep}}/K)$ is given by the cocycle

$$\phi^{\sharp} : \mathrm{Gal}(K^{\mathrm{sep}}/K) \to E(K^{\mathrm{sep}}), \quad g \mapsto g(\frac{P_c}{l^n}) - \frac{P_c}{l^n}$$

which need not necessarily be annihilated by l^n.

For any element $g \in \mathrm{Gal}(K^{\mathrm{sep}}/K)$, denote by

$$\frac{(g-1)P_c}{l^n}$$

the unique l^nth root of $(g-1)P_c$ in $E(K[c])$. This root exists because P_c belongs to $({}_{l^n}E(K[c])^{\mathcal{G}_c}$; furthermore, it is unique because $E(K[c])_{l^\infty} = 0$ (by Definition 3.1.3(f) and Proposition 1.10.1, [1, Lemma 7.14.11(i)].

The cochain

$$\psi : \mathrm{Gal}(K^{\mathrm{sep}}/K) \to E(K[c]), \quad g \mapsto -\frac{(g-1)P_c}{l^n},$$

is a cocycle whose restriction to the subgroup $\mathrm{Gal}(K^{\mathrm{sep}}/K[c])$ is the zero cochain. But ψ need not be annihilated by l^n. The cochain

$$\phi^{\sharp} + \psi : \mathrm{Gal}(K^{\mathrm{sep}}/K) \to E(K^{\mathrm{sep}}), g \mapsto g(\frac{P_c}{l^n}) - \frac{P_c}{l^n} - \frac{(g-1)P_c}{l^n},$$

is a cocycle which is annihilated by l^n and whose restriction to $\mathrm{Gal}(K^{\mathrm{sep}}/K[c])$ is the cocycle

$$\phi : \mathrm{Gal}(K^{\mathrm{sep}}/K[c]) \to E_{l^n}(K^{\mathrm{sep}}).$$

Hence the cochain $\phi^{\sharp} + \psi$ is a cocycle

$$\phi^{\sharp} + \psi : \mathrm{Gal}(K^{\mathrm{sep}}/K) \to E_{l^n}(K^{\mathrm{sep}})$$

and this cochain represents the cohomology class $\gamma_n(c)$ in $H^1(K, E_{l^n})$. Therefore the cohomology class $\delta_n(c)$ of $H^1(K[c]/K, E)_{l^n}$ is represented by the cocycle

$$\psi : \mathrm{Gal}(K[c]/K) \to E(K^{\mathrm{sep}}), \quad g \mapsto -\frac{(g-1)P_c}{l^n}$$

as required.

(iii) This follows immediately from the explicit cocycle formulae of parts (i) and (ii) of this lemma. □

4.2.2. Lemma. *Denote by a superscript* ± 1 *the eigenspaces under the action of the non-trivial element of* $\mathrm{Gal}(K/F)$. *If* $c \in \Lambda^r(n)$ *then we have*

$$P_c \pmod{l^n} \ \in \left(({}_{l^n}E(K[c]))^{\mathcal{G}_c} \right)^{-\nu(c)}$$

$$\gamma_n(c) \in H^1(K, E_{l^n})^{-\nu(c)}$$

$$\delta_n(c) \in H^1(K[c]/K, E)_{l^n}^{-\nu(c)}$$

where $\mathcal{G}_c = Gal(K[c]/K)$,

$$\nu(c) = (-1)^r \epsilon,$$

r is the number of distinct prime divisors in the support of c, and ϵ is the sign in the functional equation of the L-function of E/F.

[For the proof, see [1, Lemma 7.14.11]. Note that the set of prime numbers $\mathcal{P}$ is contained in the set of prime numbers denoted $\mathcal{P} \setminus \mathcal{F}$ of [1, Lemma 7.14.11].] □

4.3. Points P_c defined over local fields

The notation and hypotheses of (4.1.1) hold in this section. The Drinfeld-Heegner points (a, I_1, c), (a, I_1, c, π) are defined in (3.4.4) and (3.4.5).

4.3.1. Proposition. *Suppose that $c \in \Lambda(1)$ and $z \in \Lambda^1(1)$ is a prime divisor disjoint from the support of the divisor c. Let y be the unique prime of K lying over z and K_y be the completion of K at y. Then the point $(a, I_1, c) \in X_0^{\mathrm{Drin}}(K[c])$ is definable over K_y that is to say*

$$(a, I_1, c) \in X_0^{\mathrm{Drin}}(K_y).$$

Furthermore, we have that

$$(a, I_1, c, \pi) \in E(K_y)$$

is a point of the elliptic curve E definable over K_y.

Proof of Proposition 4.3.1. The prime z is inert and unramified in K/F by Definition 3.2.4 and remark 3.2.5(i). Furthermore the elliptic curve E/F has good reduction at z by remark 3.2.5(i). As I is the conductor of E/F without the component at ∞, by (4.1.1), we have that z is disjoint from the support of I and hence the curve $X_0^{\mathrm{Drin}}(I)/F$ also has good reduction z where there may be several disoint components in the closed fibre over z.

By [1, Theorem 4.6.19(ii)] as z is inert and unramified in K/F the reduction (a, I_1, c) mod z is defined over the quadratic extension field $\kappa(y)$ of $\kappa(z)$. That is to say (a, I_1, c) mod z is a point of the reduction at z of $X_0^{\mathrm{Drin}}(I)$ which is defined over $\kappa(y)$.

Let F_z be the completion of F at z. As $X_0^{\mathrm{Drin}}(I)$ has good reduction at z, it follows that the point (a, I_1, c) is defined over the field K_y as this is the unique quadratic extension field of the local field F_z which is unramified over z. It then

immediately follows that (a, I_1, c, π), which is the image of (a, I_1, c) under the morphism $\pi : X_0^{\mathrm{Drin}} \to E$ of F-schemes (see (3.4.4) and (3.4.5)), is a point of the elliptic curve E defined over K_y. □

4.3.2. Proposition. *If $z \in \Lambda^1(1)$ is a prime divisor disjoint from the support of the divisor $c \in \Lambda(1)$ and y is the prime of K lying over z, then the image of P_c in ${}_{l^n}E(K_y)$ via Proposition 4.3.1 is uniquely determined by P_c.*

Proof of Proposition 4.3.2. We have the isomorphism

$$K[c] \otimes K_y \cong \prod_i K[c]_{x_i}$$

where x_i are the places of $K[c]$ over the place y of K and $K[c]_{x_i}$ is the completion of $K[c]$ at x_i. The galois group $\mathcal{G}_c = \mathrm{Gal}(K[c]/K)$ permutes transitively the places x_i and the completions $K[c]_{x_i}$. We then have

$${}_{l^n}E(K[c] \otimes_K K_y) \cong \prod_i {}_{l^n}E(K[c]_{x_i}).$$

The point P_c belongs to $E(K[c])$ by construction. Hence the point $P_c \pmod{l^n}$ of ${}_{l^n}E(K[c])$ induces an element $(q_1, q_2, \ldots)$ of ${}_{l^n}E(K[c] \otimes_K K_y)$ where

$$(q_1, q_2, \ldots) \in \prod_i {}_{l^n}E(K[c]_{x_i})$$

and

$$q_i \in {}_{l^n}E(K[c]_{x_i}) \quad \text{for all } i.$$

By Proposition 4.3.1, P_c is definable over K_y that is to say $P_c \in E(K_y)$ and so we have $q_i \in {}_{l^n}E(K_y)$ for all i. But $P_c \pmod{l^n} \in \left({}_{l^n}E(K[c])\right)^{\mathcal{G}_c}$ by (3.4.13) (see also Lemma 4.2.2). It follows that $(q_1, q_2, \ldots)$ is invariant under $\mathcal{G}_c$. But the elements $q_i \in {}_{l^n}E(K_y)$ are permuted transitively by $\mathcal{G}_c$. Hence the elements q_i are all equal and $P_c \pmod{l^n}$ is the point $(q_1, q_1, \ldots) \in \prod_i {}_{l^n}E([c]_{x_i})$ which is in the image of the diagonal map ${}_{l^n}E(K_y) \to \prod_i {}_{l^n}E(K[c]_{x_i})$. Hence the components of the point $P_c \pmod{l^n}$ in ${}_{l^n}E(K[c] \otimes_K K_y)$ are independent of the place x_i and depend only on P_c as required. □

4.4. The map χ_z

The notation and hypotheses of (4.1.1) hold in this section.

4.4.1. Proposition. *Let $z \in \Lambda^1(n)$ be a prime divisor and let $\mathcal{E}_0/\kappa(z)$ be the closed fibre of the Néron model of E/F at the place z. Let $a_z \in \mathbb{Z}$ be the trace*

of the Frobenius at z on the inertia invariant part of the Tate module of E as in (3.2.6). Let y be the unique place of K over z. Let $c \in \Lambda(n)$ be a divisor whose support contains z and put $c' = c - z \in \Lambda(n)$ so that c' has support coprime to z.
(i) *The endomorphism $|G(c/c')|\mathrm{Frob}(z) - a_z$ of the elliptic curve $\mathcal{E}_0/\kappa(z)$ annihilates the abelian group $\mathcal{E}_0(\kappa(y))$.*
(ii) *The group homomorphism*

$$h : {}_{l^n}\mathcal{E}_0(\kappa(y)) \ \to \ \mathcal{E}_0(\kappa(y))_{l^n}, \quad x \mapsto \Big(\frac{|G(c/c')|\mathrm{Frob}(z) - a_z}{l^n}\Big)x$$

is an isomorphism which commutes with τ.

Proof of Proposition 4.4.1. (i) As in §1.4, $G(c/c')$ denotes the galois group $\mathrm{Gal}(K[c]/K[c'])$. The Frobenius $\mathrm{Frob}(z)$ acts on the Tate module of the elliptic curve $\mathcal{E}_0/\kappa(z)$ with characteristic polynomial

$$X^2 - Xa_z + |\kappa(z)|.$$

Writing F for $\mathrm{Frob}(z)$ it follows that $F^2 - Fa_z + |\kappa(z)|$ annihilates the abelian group $\mathcal{E}_0(\kappa(y))$. But F^2 is the identity automorphism on $\mathcal{E}_0(\kappa(y))$ as $\kappa(y)/\kappa(z)$ is a quadratic extension of finite fields. Hence $F^2 - Fa_z + F^2|\kappa(z)|$ annihilates $\mathcal{E}_0(\kappa(y))$. That is to say $F(F(1 + |\kappa(z)|) - a_z)$ annihilates $\mathcal{E}_0(\kappa(y))$. As F is an automorphism of $\mathcal{E}_0(\overline{\kappa(y)})$ we obtain that $F(1+|\kappa(z)|)-a_z$ annihilates $\mathcal{E}_0(\kappa(y))$.

As $K \neq F \otimes_{\mathbb{F}_q} \mathbb{F}_{q^2}$ (by the hypothesis (c) of paragraph (4.1.1)) we have that the unit group B^*/A^* is the trivial group. The Galois group

$$G(c/c') = \mathrm{Gal}(K[c]/K[c'])$$

therefore has order (see [1, Chap. 2, (2.3.8) and (2.3.12)])

$$|G(c/c')| = |\kappa(z)| + 1.$$

Hence the endomorphism $|G(c/c')|\mathrm{Frob}(z)-a_z$ of $\mathcal{E}_0$ annihilates the abelian group $\mathcal{E}_0(\kappa(y))$ as required.
(ii) Let $\alpha, \beta \in \mathbb{C}$ be the complex roots of the characteristic polynomial of Frobenius $\mathrm{Frob}(z)$ acting on the Tate module of $\mathcal{E}_0/\kappa(z)$

$$X^2 - Xa_z + |\kappa(z)|.$$

Then we have by the trace formula for the Frobenius automorphism, where $G(c/c')$ is cyclic of order $|\kappa(z)| + 1$ as in the proof of part (i),

$$|\mathcal{E}_0(\kappa(z))| = |G(c/c')| - \alpha - \beta = |G(c/c')| - a_z$$

and

$$|\mathcal{E}_0(\kappa(y))| = |\kappa(z)|^2 + 1 - \alpha^2 - \beta^2 = (|G(c/c')| - a_z)(|G(c/c')| + a_z).$$

We then obtain the decomposition into eigenspaces under the action of the involution τ, the non-trivial element of $\mathrm{Gal}(K/F)$,

$$\mathcal{E}_0(\kappa(y)) \cong \mathcal{E}_0(\kappa(z)) \oplus \mathcal{E}_0(\kappa(y))^-$$

where

$$|\mathcal{E}_0(\kappa(y))^\delta| = |G(c/c')| - \delta a_z \ \text{ for } \delta = \pm 1.$$

Hence the order of the l^∞-torsion is given by

$$|\mathcal{E}_0(\kappa(y))^\delta_{l^\infty}| = l^{s(\delta)}$$

where $l^{s(\delta)}$ is the highest power of l dividing $\delta|G(c/c')| - a_z$.

By Lemma 3.2.7(ii), we have group isomorphisms for the l^n-torsion

$$\mathcal{E}_0(\kappa(y))^\delta_{l^n} \cong \frac{\mathbb{Z}}{l^n\mathbb{Z}} \ \text{ for } \delta = \pm 1$$

that is to say these groups are cyclic of order l^n. It follows from this that there are isomorphisms for the l^∞-torsion

$$\mathcal{E}_0(\kappa(y))^\delta_{l^\infty} \cong \frac{\mathbb{Z}}{l^{s(\delta)}\mathbb{Z}} \ \text{ for } \delta = \pm 1.$$

We obtain that the l^∞-torsion subgroups of $\mathcal{E}_0(\kappa(y))^-_{l^\infty}$ and $\mathcal{E}_0(\kappa(y))^+_{l^\infty}$ are both cyclic and denoting by $|.|_l$ the normalised l-adic absolute value on $\mathbb{Q}$ we have

$$|\mathcal{E}_0(\kappa(y))^\delta_{l^\infty}| = |\ \ \delta|G(c/c')| - a_z|_l^{-1} \ \text{ for } \delta = \pm 1.$$

Let g be the homomorphism

$$g = \frac{|G(c/c')|\mathrm{Frob}(z) - a_z}{l^n} : \ \mathcal{E}_0 \to \mathcal{E}_0$$

where the integers $|G(c/c')|$, a_z are both divisible by l^n by Lemma 3.2.7(i). The homomorphism induced by g on the $\kappa(y)$-rational points of $\mathcal{E}_0$

$$g : \mathcal{E}_0(\kappa(y)) \to \mathcal{E}_0(\kappa(y))$$

is annihilated by l^n by part (i); furthermore, the subgroup $l^n\mathcal{E}_0(\kappa(y))$ of $\mathcal{E}_0(\kappa(y))$ belongs to the kernel of g again by part (i). Hence g induces a homomorphism

$$h : {}_{l^n}\mathcal{E}_0(\kappa(y)) \to \mathcal{E}_0(\kappa(y))_{l^n}.$$

On each eigencomponent $({}_{l^n}\mathcal{E}_0(\kappa(y)))^\delta$ under the action of τ, the non-trivial element of $\mathrm{Gal}(K/F)$ and where $\delta = \pm 1$, the map g is multiplication by the integer

$$N(\delta) = \frac{\delta|G(c/c')| - a_z}{l^n} = \frac{\delta|\mathcal{E}_0(\kappa(y))^\delta|}{l^n}.$$

It follows from this formula for $N(\delta)$ that the restriction of h to each eigencomponent $({}_{l^n}\mathcal{E}_0(\kappa(y)))^\delta$ is an injection. As τ commutes with h, the homomorphism h preserves the τ-eigencomponents and therefore h is an injection. As ${}_{l^n}\mathcal{E}_0(\kappa(y))$ and $\mathcal{E}_0(\kappa(y))_{l^n}$ have the same number l^{2n} of elements it follows that h is an isomorphism. □

4.4.2. Proposition. *Let z be a prime divisor of F which belongs to $\Lambda^1(n)$. Let $y \in \Sigma_K$ be the unique place of K lying over z. Assume that the prime number $l \in \mathcal{P}$ is coprime to the order of* $\mathrm{Pic}(A)$*, the Picard group of the ring A. Then there is a homomorphism*

$$\chi_z : {}_{l^n}E(K_y) \to H^1(K_y, E_{l^n})$$

with the following properties.
(1) *Let x be a place of the ring class field $K[z]$ above the place z of K. Then the image of χ_z is contained in the subgroup $H^1(K[z]_x/K_y, E_{l^n}(K[z]_x))$.*
(2) *The homomorphism χ_z is injective.*
(3) *The composition of χ_z with the homomorphism, obtained from the inclusion of group schemes $E_{l^n} \subset E$,*

$$H^1(K_y, E_{l^n}) \to H^1(K_y, E)_{l^n}$$

is an isomorphism

$${}_{l^n}E(K_y) \cong H^1(K_y, E)_{l^n}.$$

(4) *For all divisors $c \in \Lambda(n)$ such that z belongs to* $\mathrm{Supp}(c)$*, the cohomology class $\gamma_n(c)$ satisfies*

$$\gamma_n(c)_y = \chi_z(P_{c-z} \pmod{l^n})$$

where $P_{c-z} \pmod{l^n}$ is an element of ${}_{l^n}E(K_y)$ (Proposition 4.3.2) and $c - z$ has support coprime to z.

Proof of Proposition 4.4.2. For all divisors $d \in \Lambda(n)$ with support coprime to z, by Propositions 4.3.1 and 4.3.2 the point $P_d \in E(K[c])$ induces an element of ${}_{l^n}E(K_y)$ which is uniquely determined by P_d where y is the place of K over z (remark 3.2.5(1)).

Select a divisor $c \in \Lambda(n)$ with support containing z and put

$$c' = c - z \in \Lambda(n).$$

where $\mathrm{Supp}(c')$ is coprime to z.

Let z' be a prime divisor of $K[c']$ over the place y, which is the place of K over z. The prime divisor z' is totally ramified in the field extension $K[c]/K[c']$; this follows via class field theory from the definition of the ring class field $K[c]$ (see §1.4; more details are given in [1, Chapter 2, (2.3.13)]). Let $z^{\times}$ be the unique prime divisor of $K[c]$ lying over z'.

The cohomology class $\delta_n(c)$ belongs to $H^1(K[c]/K, E(K[c]))_{l^n}$ which is contained via the inflation map in $H^1(K, E(K^{\mathrm{sep}}))_{l^n}$ (diagram (3.4.12)).

Let $\mathcal{E}_0/\kappa(z)$ be the closed fibre above z of the Néron model of E/F. We then define a composite isomorphism Φ as follows where the maps i, h, j are explained below

$$\Phi : {}_{l^n}E(K_y) \xrightarrow{i} {}_{l^n}\mathcal{E}_0(\kappa(y)) \xrightarrow{h} \mathcal{E}_0(\kappa(y))_{l^n} \xrightarrow{j} E(K_y)_{l^n}. \tag{4.4.3}$$

Here $i : {}_{l^n}E(K_y) \longrightarrow {}_{l^n}\mathcal{E}_0(\kappa(y))$ is the isomorphism obtained from the surjective homomorphism $E(K_y) \longrightarrow \mathcal{E}_0(\kappa(y))$ of reduction at z whose kernel is a pro-p-group. The map $h : {}_{l^n}\mathcal{E}_0(\kappa(y)) \longrightarrow \mathcal{E}_0(\kappa(y))_{l^n}$ is the isomorphism of Proposition 4.4.1(ii). The map $j : \mathcal{E}_0(\kappa(y))_{l^n} \longrightarrow E(K_y)_{l^n}$ is the isomorphism obtained from reduction modulo y; the map j is an isomorphism because E has good reduction at y and the prime number l is distinct from the characteristic of F. The map Φ is an isomorphism as i, h, j are isomorphisms.

Let x' be a prime of $K[0]$ lying over y and let x be the unique prime of $K[z]$ over x', where y is the prime of K over z. The prime x over x' is uniquely determined by x' because $K[z]/K[0]$ is totally ramified at x' (see §1.4). The restriction of elements of the Galois group $G(c/c')$ to the fields $K[z]$ and $K[z]_x$ induces isomorphisms

$$\mathrm{Gal}(K[z]_x/K[0]_{x'}) \cong G(z/0) \cong G(c/c-z) \tag{4.4.4}$$

where these groups are cyclic of order $|\kappa(z)|+1$ (by [1, Chapter 2, (2.3.12), p.20], and as B^*/A^* is the trivial group by hypothesis (c) of (4.1.1)). Put

$$G = \mathrm{Gal}(K[z]_x/K[0]_{x'}).$$

We have the exact sequence of galois groups

$$0 \to G \to \mathrm{Gal}(K[z]_x/K_y) \to \mathrm{Gal}(K[0]_{x'}/K_y) \to 0.$$

As the field extension $K[z]/K[0]$ is totally ramified at the place x' of $K[0]$ over z, the restriction homomorphism of $G(c/0)$ to $K[0]_{x'}$ gives the isomorphism

$$G \cong G(z/0)$$

of (4.4.4).

As the field extension $K[z]_x/K[0]_{x'}$ is totally ramified, we have that the group G acts trivially on $E_{l^n}(K[z]_x)$ and that

$$E_{l^n}(K[z]_x) = E_{l^n}(K[0]_{x'}).$$

Hence we have an isomorphism

$$H^1(G, E_{l^n}(E(K[z]_x)) \cong \mathrm{Hom}(G, E_{l^n}(K[0]_{x'})). \tag{4.4.5}$$

We have already fixed in (3.4.7) a generator σ_z of $G(c/c')$. Let $\sigma \in G$ be the generator induced by σ_z under the isomorphism $G(c/c') \cong G$ of (4.4.4).

For any $P \in {}_{l^n}E(K_y)$, define a homomorphism

$$f_P : G \to E_{l^n}(K_y) \tag{4.4.6}$$

as follows. Put, where σ is the chosen generator of the cyclic group G,

$$f_P(\sigma) = \Phi(P).$$

As $\Phi(P)$ is a point of $E_{l^n}(K_y)$ and as the order of the cyclic group G is equal to $|\kappa(z)|+1$ which is divisible by l^n, the homomorphism f_P is well defined. Hence f_P defines a cohomology class in $H^1(G, E_{l^n}(K_y))$ where G acts trivially on $E_{l^n}(K_y)$.

As $\Phi : {}_{l^n}E(K_y) \to E(K_y)_{l^n}$ is an isomorphism (see (4.4.3)) and G is cyclic of order divisible by l^n, this map $P \mapsto f_P$ defines a group isomorphism

$$f : {}_{l^n}E(K_y) \xrightarrow{\cong} H^1(G, E_{l^n}(K_y)), \quad P \mapsto f_P. \tag{4.4.7}$$

We have the Hochschild-Serre spectral sequence

$$E_2^{i,j} \Rightarrow H^{i+j}(K[c]_{z^\times}/K_y, E_{l^n}(K[c]_{z^\times}))$$

where we write

$$G_{z^\times/z'} = \mathrm{Gal}(K[c]_{z^\times}/K[c']_{z'}).$$

and

$$E_2^{i,j} = H^i(K[c']_{z'}/K_y, H^j(G_{z^\times/z'}, E_{l^n}(K[c]_{z^\times}))).$$

The short exact sequence of low degree terms attached to this spectral sequence in part takes the form

$$0 \to E_2^{1,0} \to H^1(K[c]_{z^\times}/K_y, E_{l^n}(K[c]_{z^\times})) \to E_2^{0,1} \to E_2^{2,0}.$$

As the order of $\mathrm{Pic}(A)$ is coprime to l we have that the order of the place y above z in $\mathrm{Pic}(B)$ is coprime to l, as y is the unique place of K over z. It follows that the degree of the field extension $K[c']_{z'}/K_y$, which is equal to the degree of the residue field extensions, is coprime to l. Hence we have

$$H^i(K[c']_{z'}/K_y, H^j(G_{z^\times/z'}, E_{l^n}(K[c]_{z^\times}))) \cong 0 \ \text{ for all } i \geq 1$$

as the group $H^j(G_{z^\times/z'}, E_{l^n}(K[c]_{z^\times}))$ is l-power torsion for all i. The above short exact sequence of low degree terms then becomes an isomorphism
(4.4.8)

$$H^1(K[c]_{z^\times}/K_y, E_{l^n}(K[c]_{z^\times})) \cong H^0(K[c']_{z'}/K_y, H^1(G_{z^\times/z'}, E_{l^n}(K[c]_{z^\times}))).$$

As the field extension $K[c]_{z^\times}/K[c']_{z'}$ is totally ramified, we have $E_{l^n}(K[c]_{z^\times}) = E_{l^n}(K[c']_{z'})$ and the group $G_{z^\times/z'}$ acts trivially on $E_{l^n}(K[c]_{z^\times})$. This last isomorphism of (4.4.8) then becomes the isomorphism

$$H^1(K[c]_{z^\times}/K_y, E_{l^n}(K[c]_{z^\times})) \cong H^0(K[c']_{z'}/K_y, \mathrm{Hom}(G_{z^\times/z'}, E_{l^n}(K[c']_{z'})))$$

where $G_{z^\times/z'}$ acts trivially on $E_{l^n}(K[c']_{z'})$. This then provides the isomorphism

$$H^0(K[c']_{z'}/K_y, H^1(G_{z^\times/z'}, E_{l^n}(K[c]_{z^\times}))) \cong \mathrm{Hom}(G_{z^\times/z'}, E_{l^n}(K_y)).$$

This isomorphism combined with the isomorphism of (4.4.8) gives the isomorphism, where $G_{z^\times/z'}$ acts trivially on $E_{l^n}(K_y)$,

$$H^1(K[c]_{z^\times}/K_y, E_{l^n}(K[c]_{z^\times})) \cong H^1(G_{z^\times/z'}, E_{l^n}(K_y)). \tag{4.4.9}$$

Take now $c = z$ in the isomorphism of (4.4.9) where x' is a place of $K[0]$ over y and x is the unique place of $K[z]$ over x'; we have that the composition of this isomorphism (4.4.9) with the isomorphism f of (4.4.7) gives the isomorphism

$$\phi : {}_{l^n}E(K_y) \xrightarrow{\cong} H^1(K[z]_x/K_y, E_{l^n}(K[z]_x)). \tag{4.4.10}$$

The inflation map from $\mathrm{Gal}(K[z]_x/K_y)$ to $\mathrm{Gal}(K_y^{\mathrm{sep}}/K_y)$ is an injective homomorphism

$$H^1(K[z]_x/K_y, E_{l^n}(K[z]_x)) \xrightarrow{\mathrm{inf}} H^1(K_y, E_{l^n}(K_y^{\mathrm{sep}})). \tag{4.4.11}$$

The composite of this injective inflation map of (4.4.11) with the isomorphism ϕ of (4.4.10) is then defined to be the injective homomorphism χ_z

$$\begin{array}{rccc} \chi_z : & {}_{l^n}E(K_y) & \to & H^1(K_y, E_{l^n}) \\ & P & \mapsto & \{\sigma \mapsto \Phi(P)\}. \end{array} \tag{4.4.12}$$

Here $\{\sigma \mapsto \Phi(P)\}$ denotes the cohomology class f_P of (4.4.6) and (4.4.7) defined by $\sigma \mapsto \Phi(P)$ where $\sigma \in G$ is the chosen generator of G; this f_P then defines a cohomology class in $H^1(K[z]_x/K_y, E_{l^n}(K[z]_x))$ by the isomorphism of (4.4.9) for $c = z$. By inflation, this f_P gives a cohomology class of $\mathrm{Gal}(K_y^{\mathrm{sep}}/K_y)$ and hence an element of the cohomology group $H^1(K_y, E_{l^n})$ and this defines the homomorphism χ_z. This map χ_z of (4.4.12) is injective by construction.

To prove property (1), by construction the homomorphism

$$\chi_z : {}_{l^n}E(K_y) \longrightarrow H^1(K_y, E_{l^n})$$

takes a point $P \in {}_{l^n}E(K_y)$ to an element f_P of $H^1(G, E_{l^n}(K[z]_x))$ which is inflated to an element of $H^1(K_y, E_{l^n}(K^{\mathrm{sep}}))$ and this proves the property (1) of the map χ_z.

To prove property (2), the map

$$\chi_z : {}_{l^n}E(K_y) \to H^1(K_y, E_{l^n}), P \mapsto \{\sigma \mapsto \Phi(P)\}$$

is injective by construction, as already noted.

For property (3), we have by definition that the map χ_z of (4.4.12) is the composition of the isomorphism ϕ of (4.4.10) with the inflation map of (4.4.11). That is to say, the map χ_z factors as

$$(4.4.13) \qquad {}_{l^n}E(K_y) \xrightarrow{\phi} H^1(K[z]_x/K_y, E_{l^n}(K[z]_x)) \xrightarrow{\mathrm{inf}} H^1(K_y, E_{l^n}(K_y^{\mathrm{sep}}))$$

where ϕ is an isomorphism and inf is an injection.

We have an exact sequence obtained from the inclusion of group schemes $E_{l^n} \subset E$,

$$0 \longrightarrow {}_{l^n}E(K_y) \xrightarrow{\partial_{l^n}} H^1(K_y, E_{l^n}) \xrightarrow{\psi} H^1(K_y, E)_{l^n}.$$

The morphism of multiplication by l^n on the Néron model of E over the ring of valuation integers of F at the place z is étale; therefore the image of ∂_{l^n} consists of unramified cohomology classes and more precisely the image of ∂_{l^n} belongs to the subgroup $H^1(K_y^{\mathrm{nr}}/K_y, E_{l^n})$ of $H^1(K_y, E_{l^n})$ where K_y^{nr} is the maximal separable unramified extension of K_y. The intersection of $H^1(K_y^{\mathrm{nr}}/K_y, E_{l^n})$ with $H^1(K[z]_x/K_y, E_{l^n}(K[z]_x))$, which are both subgroups of $H^1(K_y, E_{l^n})$, is therefore contained in $H^1(K[0]_{x'}/K_y, E_{l^n}(K[0]_{x'}))$; this follows by considering representative cocycles and because the field extension $K[z]_x/K[0]_{x'}$ is totally ramified and $K[0]_{x'}/K_y$ is unramified. But the order of the Picard group $\mathrm{Pic}(A)$ is coprime to l by hypothesis; therefore the order of the place y over z in $\mathrm{Pic}(B)$ is coprime to l as y is the unique place of K over z. It follows that the degree of the field extension $K[0]_{x'}/K_y$ is coprime to l and therefore we have

$$H^1(K[0]_{x'}/K_y, E_{l^n}(K[0]_{x'})) = 0.$$

It follows that the composition of the injective inflation map

$$H^1(K[z]_x/K_y, E_{l^n}(K[z]_x)) \xrightarrow{\mathrm{inf}} H^1(K_y, E_{l^n}(K_y^{\mathrm{sep}}))$$

with $\psi : H^1(K_y, E_{l^n}) \to H^1(K_y, E)_{l^n}$ is an injection

$$(4.4.14) \qquad H^1(K[z]_x/K_y, E_{l^n}(K[z]_x)) \longrightarrow H^1(K_y, E)_{l^n}.$$

By arithmetic flat local duality, there is an isomorphism of discrete groups

$$(4.4.15) \qquad H^1(K_y, E) \cong \mathrm{Hom}(E(K_y), \mathbb{Q}/\mathbb{Z})$$

for the proof of which, using flat cohomology and local class field theory, see [9, Chapter 3, Theorem 7.8]; we only require the prime-to-p version of this duality (4.4.15) and whose simpler proof is explained in [9, Chapter 1, Remark 3.6]. From (4.4.15) we obtain the isomorphism of discrete groups

$$H^1(K_y, E)_{l^n} \cong \mathrm{Hom}({}_{l^n}E(K_y), \mathbb{Z}/l^n\mathbb{Z}). \tag{4.4.16}$$

It follows from (4.4.16) that $H^1(K_y, E)_{l^n}$ and ${}_{l^n}E(K_y)$ have the same number of elements. As $E_{l^n}(K_y)$ is isomorphic to ${}_{l^n}E(K_y)$ by the isomorphism Φ of (4.4.3), we have that $H^1(K_y, E)_{l^n}$ and $E_{l^n}(K_y)$ have the same number of elements.

From (4.4.9) for $c = z$, we have the isomorphism, where G is cyclic of order divisible by l^n and which acts trivially on $E_{l^n}(K_y)$,

$$H^1(K[z]_x/K_y, E_{l^n}(K[z]_x)) \cong H^1(G, E_{l^n}(K_y)) \cong E_{l^n}(K_y).$$

It now follows that the finite groups $H^1(K[z]_x/K_y, E_{l^n}(K[z]_x))$, $E_{l^n}(K_y)$ and $H^1(K_y, E)_{l^n}$ have the same number of elements and hence the injective homomorphism of (4.4.14)

$$H^1(K[z]_x/K_y, E_{l^n}(K[z]_x)) \longrightarrow H^1(K_y, E)_{l^n}$$

is an isomorphism. It follows that the natural homomorphism, obtained from the inclusion of group schemes $E_{l^n} \subset E$,

$$H^1(K_y, E_{l^n}) \xrightarrow{\psi} H^1(K_y, E)_{l^n}$$

composed with χ_z, where χ_z factors through the group $H^1(K[z]_x/K_y, E_{l^n}(K[z]_x))$ as in (4.4.13),

$${}_{l^n}E(K_y) \xrightarrow{\chi_z} H^1(K_y, E_{l^n}) \xrightarrow{\psi} H^1(K_y, E)_{l^n}$$

is an isomorphism

$${}_{l^n}E(K_y) \cong H^1(K_y, E)_{l^n}.$$

This proves the property (3).

It only remains to prove property (4). For all divisors $d \in \Lambda(n)$, as in (3.4.10) we shall write $P_d \pmod{l^n}$ for the image of $P_d \in E(K[d])$ in ${}_{l^n}E(K[d])$.

From Lemma 4.2.1, we have that the cohomology class $\gamma_n(c)$ in $H^1(K, E_{l^n})$ is represented by the cocycle, where $\frac{(\sigma-1)P_c}{l^n}$ is the unique l^n-division point of $(\sigma - 1)P_c$ in $E(K[c])$ and P_c/l^n is a fixed l^n-division point of P_c,

$$\Gamma_n(c) : \sigma \mapsto -\frac{(\sigma-1)P_c}{l^n} + \sigma\frac{P_c}{l^n} - \frac{P_c}{l^n}, \quad \mathrm{Gal}(K^{\mathrm{sep}}/K) \to E(K^{\mathrm{sep}})_{l^n} \tag{4.4.17}$$

and the cohomology class $\delta_n(c)$ in $H^1(K, E)_{l^n}$ is represented by the cocycle

$$\Delta_n(c) : \sigma \mapsto -\frac{(\sigma-1)P_c}{l^n}, \quad \mathrm{Gal}(K^{\mathrm{sep}}/K) \to E(K^{\mathrm{sep}}). \tag{4.4.18}$$

The cohomology class $\delta_n(c)_y \in H^1(K_y, E(K_y^{\mathrm{sep}}))$ is the restriction at y of the class $\delta_n(c) \in H^1(K, E(K^{\mathrm{sep}}))$. Let $Q \in E(K[c])$ be the element given by

$$Q = -\frac{(\sigma_z - 1)P_c}{l^n} \tag{4.4.19}$$

and where σ_z is the fixed generator of the cyclic group $G(c/c')$.

The definition of P_c (as in [1, Chap. 7, §7.14.10 pp.386-387] and (3.4.9) above) is the following. Let $c = \sum_{i=1}^{r} z_i$ be the decomposition of c as a sum of distinct prime divisors, where we write $z = z_1$. Then we have

$$P_c = \sum_{s \in \mathcal{S}} s D_c y_c$$

where

$$y_c = (0, I_1, c, \pi) \in E(K[c]) \tag{4.4.20}$$

as in (3.4.7) and where $\mathcal{S}$ is a set of coset representatives for $G(c/0)$ in $\mathrm{Gal}(K[c]/K)$ and D_c is the Kolyvagin element of (3.4.9).

We have an exact sequence of abelian groups

$$0 \to G(c/c') \to G(c/0) \to G(c'/0) \to 0.$$

We obtain a corresponding decomposition in the group algebra $\mathbb{Z}[G(c/0)]$ of the Kolyvagin element D_c

$$D_c = D_{c-z} D_z$$

We have already selected in (3.4.7) a generator σ_z of the cyclic group $G(c/c-z)$. We may then define a map of sets

$$h_z : G(c/c-z) \to \mathbb{Z}$$

by

$$\sigma_z^{-s} \mapsto -s, \quad \text{for } s = 0, 1, \ldots, |G(c/c-z)| - 1.$$

The Kolyvagin element of h_z is then as in (3.4.8)

$$D_z = -\sum_{r=1}^{|G(c/c-z)|-1} r\sigma_z^r.$$

We have

$$(\sigma_z - 1)D_z = -|G(c/c-z)| + e_{G(c/c-z)} \tag{4.4.21}$$

where $e_{G(c/c-z)}$ is the element of the group algebra $\mathbb{Z}[G(c/c-z)]$ given by

$$e_{G(c/c-z)} = \sum_{g \in G(c/c-z)} g.$$

The element $P_c \in E(K[c])$ may then be written as, where we write $y_c = (0, I_1, c, \pi)$ as in (3.4.7) and (4.4.20),

$$P_c = \sum_{s\in\mathcal{S}} sD_{c-z}D_z y_c. \tag{4.4.22}$$

We have (see [1, Chap. 4, (4.8.3) and table 4.8.5])

$$\frac{|O^*_{c'}|}{|A^*|}\mathrm{Tr}_{K[c]/K[c']}y_c = a_z y_{c'}$$

where $a_z \in \mathbb{Z}$ is as in Proposition 4.4.1 and (3.2.6) and where $y_{c'} = (0, I_1, c', \pi)$. By definition $Q = -((\sigma_z - 1)P_c)/l^n$ (see (4.4.19)); hence we have from (4.4.21)

$$Q = -\sum_{s\in\mathcal{S}} sD_{c-z}(\frac{(\sigma_z - 1)D_z}{l^n})y_c$$

$$= \sum_{s\in\mathcal{S}} sD_{c-z}(\frac{|G(c/c')|}{l^n}y_c - \frac{|A^*|}{|O^*_{c'}|}\frac{a_z}{l^n}y_{c'}).$$

As K is not obtained from F by ground field extension (hypothesis (c) of paragraph (4.1.1)), we have $A^* = O^*_{c'}$; hence we obtain

$$Q = \sum_{s\in\mathcal{S}} sD_{c-z}(\frac{|G(c/c')|}{l^n}y_c - \frac{a_z}{l^n}y_{c'}). \tag{4.4.23}$$

As at the beginning of this proof, let y be the unique place of K over z, z' be a prime of $K[c']$ over the place y of K, and let $z^\times$ be the place of $K[c]$ over the prime z' of $K[c']$ where the field extension $K[c]/K[c']$ is totally ramified at z'. Also $\mathcal{E}_0/\kappa(z)$ denotes the closed fibre above z of the Néron model of E/F.

We write Q_0 for the image of Q modulo $z^\times$ in $\mathcal{E}_0(\kappa(z^\times))$ by passage to the residue field $\kappa(z^\times)$. From the isomorphism (4.4.9), where $\delta_n(c)_y$ belongs to $H^1(K[c]_{z^\times}/K_y, E_{l^n}(K[c]_{z^\times}))$, we have that the reduction of $\delta_n(c)_y$ at y belongs to $\mathrm{Hom}(G(c/c'), \mathcal{E}_0(\kappa(y)))_{l^n}$ and is given by the cocycle (see (4.4.18))

$$g \mapsto -\frac{(g-1)P_c}{l^n} \pmod{z^\times}, \quad \mathrm{Gal}(K[c]_{z^\times}/K[c']_{z'}) \cong G(c/c') \to \mathcal{E}_0(\kappa(y))$$

and we have that $-\frac{(g-1)P_c}{l^n}$ modulo $z^\times$ belongs to the l^n-torsion subgroup $\mathcal{E}_0(\kappa(y))_{l^n}$ rational over $\kappa(y)$. Hence the point Q_0, the reduction of Q modulo $z^\times$, belongs to $\mathcal{E}_0(\kappa(y))_{l^n}$. Note that $-(g-1)P_c$ modulo $z^\times$ reduces to zero, for all $g \in G(c/c')$, as $K[c]/K[c']$ is totally ramified at z'.

Denote by $\mathrm{Frob}(z)$ the Frobenius automorphism $x \mapsto x^{|\kappa(z)|}$ of the closed fibre $\mathcal{E}_0/\kappa(z)$ over z of the Néron model of E/F. The theorem [1, Chap. 4, Theorem 4.8.9] gives that for the prime z' of $K[c']$ above z we have, where $y_c = (0, I_1, c, \pi)$ as in (4.4.20),

$$\mathrm{Frob}(z)y_c \equiv y_{c'} \pmod{z'}. \tag{4.4.24}$$

We obtain from Proposition 4.3.1 that $y_c \bmod z^\times$ is defined over the subfield $\kappa(y)$ of $\kappa(z^\times)$ where $\kappa(y)$ is the quadratic extension field of the finite field $\kappa(z)$. Hence we have from (4.4.24)

$$\frac{|G(c/c')|}{l^n}y_c - \frac{a_z}{l^n}y_{c'} \equiv \frac{|G(c/c')|\mathrm{Frob}(z) - a_z}{l^n}y_{c'} \pmod{z'}. \tag{4.4.25}$$

The point $P_{c'} \in E(K[c'])$ is given by (see (3.4.9))

$$P_{c'} = \sum_{s\in\mathcal{S}} sD_{c-z}y_{c'}.$$

We then have from (4.4.23) and (4.4.25)

$$Q = \sum_{s\in\mathcal{S}} sD_{c-z}\Big(\frac{|G(c/c')|}{l^n}y_c - \frac{a_z}{l^n}y_{c'}\Big)$$

$$\equiv\ Q_0\ \equiv \frac{|G(c/c')|\mathrm{Frob}(z) - a_z}{l^n}P_{c'}\ \pmod{z'}. \tag{4.4.26}$$

From Lemma 4.2.2 or [1, Lemma 7.14.11(ii)]), we have that $P_{c'} \pmod{l^n}$ belongs to the $-\nu(c')$-eigenspace for τ on ${}_{l^n}E(K[c'])$, where $\nu(c') = (-1)^r\epsilon$ and r is the number of prime divisors in the support of c' and ϵ is the sign in the functional equation of the L-function of E/F (as in Lemma 4.2.2). As z is inert in K/F, it follows that the image of the reduction $P_{c'}^\flat$ of $P_{c'}$ modulo z' belongs to the $-\nu(c')$-eigenspace for τ on ${}_{l^n}\mathcal{E}_0(\kappa(z'))$. Let

$$h : {}_{l^n}\mathcal{E}_0(\kappa(y)) \ \to\ \mathcal{E}_0(\kappa(y))_{l^n},\quad x \mapsto \Big(\frac{|G(c/c')|\mathrm{Frob}(z) - a_z}{l^n}\Big)x$$

be the isomorphism which commutes with τ of Proposition 4.4.1. As Q_0 belongs to the subgroup $\mathcal{E}_0(\kappa(y))$ as already noted and also as $P_{c'}^\flat \pmod{l^n}$ is an element of ${}_{l^n}\mathcal{E}_0(\kappa(y))$ by Proposition 4.3.2, we have by (4.4.26)

$$Q_0 = h(P_{c'}^\flat \pmod{l^n}).$$

With the notation of (4.4.3) we then have

$$Q_0 = h \circ i(P_{c'} \pmod{l^n})$$

where i is the reduction isomorphism ${}_{l^n}E(K_y) \to {}_{l^n}\mathcal{E}_0(\kappa(y))$ and where $P_{c'} \pmod{l^n}$ belongs to ${}_{l^n}E(K_y)$ by Proposition 4.3.2. Furthermore, by (4.4.3) and as $\Phi = j \circ h \circ i$ we have

$$\Phi(P_{c'} \pmod{l^n}) = j(Q_0) \tag{4.4.27}$$

is the unique l^n-torsion point of $E(K_y)$ whose reduction at y is Q_0.

Let $\overline{z}$ be any prime of K^{sep} above $z^{\times}$. Restrict the cocycle

$$\Gamma_n(c) : \text{Gal}(K^{\text{sep}}/K) \to E(K^{\text{sep}})_{l^n}$$

of (4.4.17), which represents $\gamma_n(c)$, to the decomposition group of $\overline{z}$. As $K[c]/K[c']$ is totally ramified at z' we have that the Kolyvagin element D_z restricted to the the residue field of $z^{\times}$ satisfies

$$D_z = -|\kappa(z)|(|\kappa(z)| + 1)/2.$$

Furthermore, l^n divides $|\kappa(z)| + 1$ and l is different from 2; hence we have that (from (4.4.22)) the reduction $P_c^{\flat}$ of P_c at $z^{\times}$ satisfies $P_c^{\flat} \in l^n \mathcal{E}_0(\kappa(z^{\times}))$ and hence as E has good reduction at $z^{\times}$ we have that

$$P_c \in l^n E(K[c]_{z^{\times}}).$$

We then have that the cocycle $\Gamma_n(c)$ of (4.4.17) satisfies

$$\Gamma_n(c)(\rho) = 0 \quad \text{for all } \rho \in \text{Gal}(K[c]_{z^{\times}}^{\text{sep}}/K[c]_{z^{\times}})$$

and this cocycle, restricted to the decomposition group of $\overline{z}$, factors through the subgroup $\text{Gal}(K[c]_{z^{\times}}/K_y)$. Furthermore, since σ_z, which is the selected generator of $G(c/c')$, is in the inertia group of y we have that

$$\sigma_z \frac{P_c}{l^n} - \frac{P_c}{l^n}$$

reduces to zero modulo $z^{\times}$. Hence

$$\Gamma_n(c)(\sigma_z) = -\frac{(\sigma_z - 1)P_c}{l^n} + \sigma_z \frac{P_c}{l^n} - \frac{P_c}{l^n}$$

is the unique l^n-torsion point congruent to $Q \pmod{z^{\times}}$ where Q is given by (4.4.19) that is to say

$$\Gamma_n(c)(\sigma_z) \equiv -\frac{(\sigma_z - 1)P_c}{l^n} \pmod{z^{\times}}. \tag{4.4.28}$$

But from (4.4.27), we then obtain

$$\Gamma_n(c)(\sigma_z) = \Phi(P_{c'} \pmod{l^n}). \tag{4.4.29}$$

Then (4.4.29) shows, as $\Gamma_n(c)(\sigma_z)$ is the unique l^n-torsion point with reduction at y coinciding with the reduction of $Q = -\frac{(\sigma_z - 1)P_c}{l^n}$ at y, that we have an equality of cohomology classes in $H^1(K_y, E_{l^n})$

$$\gamma_n(c)_y = \chi_z(P_{c-z} \pmod{l^n})$$

where $P_{c-z} \pmod{l^n}$ is an element of ${}_{l^n}E(K_y)$. This proves property (4) and completes the proof of Proposition 4.4.2. □

4.4.30. Remarks. (1) This Proposition 4.4.2 and its consequence stated in Proposition 4.5.1(iv) below are an extension of [1, Chapter 7, Lemma 7.14.14(ii)]. It would be of interest to eliminate the hypothesis that the prime number l be coprime to the order of the Picard group $\mathrm{Pic}(A)$ from these Propositions 4.4.2 and 4.5.1(iv).
(2) The homomorphism χ_z interchanges the τ-eigenspaces in that for $\delta = \pm 1$ we have

$$\chi_z({}_{l^n}E(K_y))^{\delta}) \subseteq H^1(K_y, E_{l^n})^{-\delta}.$$

This property, which is not required for this paper, follows from the group $\mathrm{Gal}(K[c]/F)$ being generalised dihedral.

4.5. Localizations of the classes $\gamma_n(c)$ and $\delta_n(c)$

The notation and hypotheses of (4.1.1) hold in this section. Let $Ш(E/K)$ be the Tate-Shafarevich group of the elliptic curve $E \times_F K$ over K.

4.5.1. Proposition. *Let $c \in \Lambda(n)$, let z be a prime divisor in $\Lambda^1(n)$ and let y be the place of K over the place z of F.*
(i) *Let v be a place of K coprime to c. Then we have $\gamma_n(c)_v \in \partial_n(E(K_v))$ that is to say we have $\delta_n(c)_v = 0$.*
(ii) *If c is coprime to z then we have $\gamma_n(c)_y = \partial_n((P_c \pmod{l^n})_y)$.*
(iii) *If $l^n | P_c$ then we have $\delta_n(c) = 0$. If $l^n | P_{c-w}$ for all prime divisors w in the support of c then we have $\delta_n(c) \in Ш(E/K)_{l^\infty}$.*
(iv) *Assume that the prime number $l \in \mathcal{P}$ is coprime to the order of* $\mathrm{Pic}(A)$*, the Picard group of the ring A. If $z \in \mathrm{Supp}(c)$ then we have*

$$\mathrm{ord}\ \delta_n(c)_y = \mathrm{ord}\ \gamma_n(c)_y = \mathrm{ord}\ \gamma_n(c-z)_y =$$

$$\mathrm{ord}\Big((P_{c-z} \pmod{l^n})_y\Big)$$

where $(P_{c-z} \pmod{l^n})_y$ is an element of ${}_{l^n}E(K_y)$.

Proof. (i) The field $K[c]$ is a subfield of K_∞ as ∞ is split completely in $K[c]/K$ (see [1, Chapter 2, (2.3.13)]). We have that the localization $\delta_n(c)_\infty$ at ∞ of $\delta_n(c)$ satisfies $\delta_n(c)_\infty \in H^1(K_\infty, E)_{l^n}$ and the localization $(P_c \pmod{l^n})_\infty$ at ∞ of $P_c \pmod{l^n}$ satisfies $(P_c \pmod{l^n})_\infty \in {}_{l^n}E(K_\infty)$. It then follows from the diagram (3.4.12) that $\delta_n(c)_\infty = 0$.

Suppose now that v is a place of K such that $v \neq \infty$ and $v \notin \mathrm{Supp}(c)$. We have by construction that

$$\delta_n(c) \in H^1(K[c]/K, E(K[c]))_{l^n}.$$

The field extension $K[c]/K$ is unramified at v (see [1, Chapter 2, (2.3.13)]). Hence we have that the localization $\delta_n(c)_v$ at v satisfies

$$\delta_n(c)_v \in H^1(K_v^{nr}/K_v, E(K_v^{nr}))_{l^n} \subseteq H^1(K_v, E(K_v^{\mathrm{sep}}))_{l^n}$$

where K_v^{nr} is the maximal unramified separable extension of the local field K_v. But $H^1(K_v^{nr}/K_v, E(K_v^{nr}))_{l^n} = 0$ by Definition 3.1.3(d) of the set of prime numbers $\mathcal{P}$ to which l belongs. Hence we have $\delta_n(c)_v = 0$; this last vanishing is equivalent to $\gamma_n(c)_v \in \partial_n(E(K_v))$, as required.
(ii) By Lemma 4.2.1, the cohomology class $\gamma_n(c)$ is represented by the cocycle

$$\sigma \mapsto -\frac{(\sigma-1)P_c}{l^n} + \sigma\frac{P_c}{l^n} - \frac{P_c}{l^n}, \quad \mathrm{Gal}(K^{\mathrm{sep}}/K) \to E(K^{\mathrm{sep}})_{l^n}$$

where $\frac{(\sigma-1)P_c}{l^n}$ is the unique l^n-division point of $(\sigma-1)P_c$ in $E(K[c])$. Furthermore by the same Lemma 4.2.1, the cohomology class $\delta_n(c) \in H^1(K, E(K^{\mathrm{sep}}))_{l^n}$ is represented by the cocycle

$$\sigma \mapsto -\frac{(\sigma-1)P_c}{l^n}, \quad \mathrm{Gal}(K^{\mathrm{sep}}/K) \to E(K^{\mathrm{sep}}).$$

By part (i) as $z \notin \mathrm{Supp}(c)$, we have $\delta_n(c)_y = 0$ hence a cocycle representing $\delta_n(c)$ when localized at v is cohomologous to zero. Therefore there is an element $a \in E(K_y^{\mathrm{sep}})$ such that

$$-\frac{(\sigma-1)P_c}{l^n} = \sigma a - a, \quad \text{for all } \sigma \in \mathrm{Gal}(K_y^{\mathrm{sep}}/K_y).$$

Hence we have

$$-(\sigma-1)(P_c + l^n a) = 0, \quad \text{for all } \sigma \in \mathrm{Gal}(K_y^{\mathrm{sep}}/K_y).$$

This implies that

$$P_c + l^n a \in E(K_y).$$

By Lemma 4.3.1, the localization at y of the point P_c lies in $E(K_y)$ where y is the place of K over z. Furthermore, by Lemma 4.3.2 the point $(P_c \pmod{l^n})_y$ belongs to ${}_{l^n}E(K_y)$ and its image in this group is uniquely determined by P_c.

It follows from this and the above cocycle formulae for $\gamma_n(c)$ and $\delta_n(c)$ that the cohomology class $\gamma_n(c)_y$ is represented by the cocyle

$$\sigma \mapsto \sigma\frac{P_c}{l^n} - \frac{P_c}{l^n}, \quad \mathrm{Gal}(K_y^{\mathrm{sep}}/K_y) \to E(K_y^{\mathrm{sep}})_{l^n}.$$

That is to say we have $\gamma_n(c)_y = \partial_n((P_c \pmod{l^n})_y)$.
(iii) From Lemma 4.2.1(ii), the cohomology class $\delta_n(c)$ is represented by the cocycle

$$\sigma \mapsto -\frac{(\sigma - 1)P_c}{l^n}, \quad \mathrm{Gal}(K^{\mathrm{sep}}/K) \to E(K^{\mathrm{sep}}).$$

We then evidently have that if $l^n | P_c$, that is to say $P_c \in l^n E(K[c])$, then $\delta_n(c) = 0$.

Suppose that $l^n | P_{c-z}$ for all prime divisors z in the support of c. Then for any prime divisor y of K lying over the prime divisor z dividing c we have $\delta_n(c)_y = 0$ by Proposition 4.4.2(4). It then follows from part (i) of the present Proposition 4.5.1 that $\delta_n(c)_v = 0$ for all places v of K and hence that $\delta_n(c)$ belongs to the Tate-Shafarevich group $Ш(E/K)_{l^\infty}$.
(iv) From the property (4) of the map χ_z of Proposition 4.4.2, we have

$$\gamma_n(c)_y = \chi_z((P_{c-z} \pmod{l^n})_y)$$

where $(P_{c-z} \pmod{l^n})_y$ is an element of ${}_{l^n}E(K_y)$ by Proposition 4.3.2. As the homomorphism χ_z is injective (property (2) of Proposition 4.4.2), we obtain

$$\mathrm{ord}(\gamma_n(c)_y) = \mathrm{ord}((P_{c-z} \pmod{l^n})_y)$$

where again $(P_{c-z} \pmod{l^n})_y$ denotes an element of ${}_{l^n}E(K_y)$.

From part (ii) of the present Proposition 4.5.1, we have

$$\gamma_n(c - z)_y = \partial_n((P_{c-z} \pmod{l^n})_y)$$

where z is coprime to the support of $c - z$. Hence we obtain

$$\mathrm{ord}((P_{c-z} \pmod{l^n})_y) = \mathrm{ord}(\gamma_n(c - z)_y).$$

From property (3) of Proposition 4.4.2, we have that the composition χ_z with the homomorphism, obtained from the inclusion of group schemes $E_{l^n} \subset E$,

$$\theta : H^1(K_y, E_{l^n}) \to H^1(K_y, E)_{l^n}$$

is an isomorphism

$${}_{l^n}E(K_y) \cong H^1(K_y, E)_{l^n}.$$

Hence the image in $H^1(K_y, E)_{l^n}$ under θ of $\chi_z(P_{c-z} \pmod{l^n})$ is the cohomology class $\delta_n(c)_y$ because this is the image under θ of $\gamma_n(c)_y \in H^1(K_y, E_{l^n})$ and furthermore, because $\theta \circ \chi_z$ is an isomorphism, $\delta_n(c)_y$ has the same order as $\mathrm{ord}((P_{c-z} \pmod{l^n})_y)$, which completes the proof of the proposition. □

4.6. The Cassels pairing with a class $\delta_n(c)$

The notation and hypotheses of (4.1.1) hold in this section. The torsion abelian group $\mathbb{Z}/n\mathbb{Z}$ for $n \neq 0$ is considered to be a subgroup of $\mathbb{Q}/\mathbb{Z}$ via the map $1 \mapsto 1/n$. Let

$Ш(E/K)$ be the Tate-Shafarevich group of the elliptic curve $E \times_F K$ over K;
$< -, - >_{\text{Cassels}}$ be the Cassels pairing on $Ш(E/K)$ (see §2.3);
$[,]_w : H^1(K_w, E)_n \times {}_nE(K_w) \to \mathbb{Z}/n\mathbb{Z}$ be the local Tate pairing for any place $w \in \Sigma_K$ of K and any integer n coprime to the characteristic of F.

4.6.1. Proposition. *Let m and n be integers ≥ 1 and let $c \in \Lambda(m+n)$. Suppose that $\delta_m(c)$ belongs to the Tate-Shafarevich group $Ш(E/K)_{l^\infty}$. Suppose that the element $s \in Ш(E/K)_{l^\infty}$ has order at most l^n. Lift the element $s \in Ш(E/K)_{l^\infty}$ to an element $S \in H^1(K, E_{l^n})$ and select points $x(w) \in E(K_w)$ such that $S_w = \partial_{l^n}(x(w))$ for all $w \in \Sigma_K$. Then we have*

$$< \delta_m(c), s >_{\text{Cassels}} = \sum_{y \in \Sigma_K \text{ divides } \text{Supp}(c)} [\delta_{m+n}(c)_y, x(y)]_y \tag{4.6.2}$$

where the sum runs over the places of K which divide an element of $\text{Supp}(c)$.

Proof. The construction of the Cassel's pairing is given in §2.3. As in the statement of the proposition, we may lift the element $s \in Ш(E/K)_{l^\infty}$ of order l^n to an element $S \in H^1(K, E_{l^n})$. The points $x(w) \in E(K_w)$ are then selected such that $S_w = \partial_{l^n}(x(w))$ for all $w \in \Sigma_K$.

By Lemma 4.2.1(iii) we have

$$l^n \delta_{m+n}(c) = \delta_m(c).$$

From the formula (2.3.5) for the Cassels pairing, we then obtain

$$< \delta_m(c), s >_{\text{Cassels}} = \sum_{w \in \Sigma_K} [\delta_{m+n}(c)_w, x(w)]_w. \tag{4.6.3}$$

where the sum runs over all places w of K.

By Proposition 4.5.1(i), we have $\delta_{m+n}(c)_w = 0$ for all places $w \in \Sigma_K$ which do not divide an element of $\text{Supp}(c)$. Hence there is no contribution to the Cassels pairing in the sum (4.6.3) when w does not divide an element of $\text{Supp}(c)$. Hence we obtain the formula (4.6.2). □

4.6.4. Proposition. *Let m and n be integers ≥ 1 and let $c \in \Lambda(m+n)$, $d \in \Lambda(n)$. Suppose that $\delta_m(c)$ and $\delta_n(d)$ belong to the Tate-Shafarevich group $Ш(E/K)_{l^\infty}$. Assume that the prime number $l \in \mathcal{P}$ is coprime to the order of*

$\mathrm{Pic}(A)$. *Then we have*

(4.6.5)

$$< \delta_m(c), \delta_n(d) >_{\mathrm{Cassels}} = \sum_{y \in \Sigma_K \text{ divides } \mathrm{Supp}(c) \setminus \mathrm{Supp}(d)} [\delta_{m+n}(c)_y, (P_d \pmod{l^n})_y]_y$$

where the sum runs over the places of K *which divide an element of* $\mathrm{Supp}(c) \setminus \mathrm{Supp}(d)$.

Proof. The element $\gamma_n(d) \in H^1(K, E_{l^n})$ is a lifting of $\delta_n(d) \in Ш(E/K)$ which has order at most l^n. Suppose that $z \in \mathrm{Supp}(c)$ and y is the unique place of K lying over the place z of F. If z also satisfies $z \in \mathrm{Supp}(d)$ then by Proposition 4.5.1(iv), we have

$$\gamma_n(d)_y = 0$$

because $\delta_n(d)_y = 0$ as we have $\delta_n(d) \in Ш(E/K)$. If on the other hand z satisfies $z \notin \mathrm{Supp}(d)$ then by Lemma 4.5.1(ii), we have

$$\gamma_n(d)_y = \partial_{l^n}((P_d \pmod{l^n})_y).$$

where $(P_d \pmod{l^n})_y$ denotes the localization at y of $P_d \pmod{l^n}$. The formula (4.6.5) to be proved now follows from the formula (4.6.2) of Proposition 4.6.1. □

Chapter 5. Construction of cohomology classes and proofs of the main theorems

5.1. M_r is finite for some r

(5.1.1) Throughout this chapter 5, the notation of (4.1.1) of the previous chapter 4 remains valid and the elliptic curve E/F and quadratic field extension K/F satisfy the hypotheses (a), (b), (c) of (4.1.1).

We further let in this chapter

N^+, N^- be the eigenspaces under the action of the involution τ whenever N is a $\mathbb{Z}[\mathrm{Gal}(K/F)]$-module on which multiplication by 2 is an isomorphism;
$\epsilon = \pm 1$ is the sign of the functional equation of the L-function $L(E/F, s)$ of E/F;
$\nu(r) = (-1)^r\epsilon$ for any natural number $r \in \mathbb{N}$;
$\nu(c) = (-1)^r\epsilon$ for any divisor c of F with exactly r distinct prime divisors in its support;
$P_c \in E(K[c])$ be the Drinfeld-Heegner points of paragraph (3.4.9) for all $c \in \Lambda(1)$, where $P_0 \in E(K)$;
M_r, for all $r \in \mathbb{N}$, be the quantities in $\mathbb{N} \cup \infty$ given in Definition 4.1.6;
$Ш(E/K)$ be the Tate-Shafarevich group of the elliptic curve $E \times_F K$ over K (see §2.2);
$\mathrm{Sel}_n(E/K)$ be the n-Selmer group of $E \times_F K$ over K for any integer n coprime to the characteristic of F (see §2.2));
$\mathrm{Sel}_{a^\infty}(E/K) = \varinjlim_n \mathrm{Sel}_{a^n}(E/K)$ for any number a coprime to the characteristic of F (see also (3.1.6)).

In this section §5.1 some consequences are presented of the hypothesis that M_r be finite for some r.

5.1.2. Lemma. *The Drinfeld-Heegner P_0 has infinite order in $E(K)$ if and only if M_0 is finite. If P_0 has infinite order then we have*

$$l^{M_0} = |(E(K)/\mathbb{Z}P_0)_{l^\infty}|$$

$$\mathrm{ord}\ \gamma_{M_0+m}(0) = l^m, \quad \textit{for all } m \geq 0,$$

$$\gamma_{M_0+m}(0) \in \mathrm{Sel}_{l^\infty}(E/K)^{-\epsilon} \quad \textit{for all } m \geq 0.$$

Proof of Lemma 5.1.2. The definition of M_0 is that (Definition 4.1.6)

$$M_0 = \mathrm{ord}_l(P_0) = \max\{m|\ P_0 \in l^m E(K[0])\}.$$

The group $E(K[0])$ has no l-torsion (as $l \in \mathcal{P}$, see Proposition 1.10.1 and Definition 3.1.3(f)) and is a finitely generated group by the Mordell-Weil theorem. Hence P_0 has infinite order in $E(K)$ if and only if M_0 is finite.

We have $\gamma_{M_0+m}(0) \in \mathrm{Sel}_{l^\infty}(E/K)^{-\epsilon}$, for all $m \geq 1$, by Lemma 4.2.2 and because $\gamma_{M_0+m}(0) = \partial_{l^{M_0+m}}(P_0)$.

Assume now that P_0 has infinite order. By [1, Chap. 7, Theorem 7.6.5] the point P_0 generates a subgroup of $E(K)$ of finite index (note that in the notation of this theorem [1, Chap. 7, Theorem 7.6.5], we have $P_0 = x_0$).

By definition we have

$$|(E(K)/\mathbb{Z}P_0)_{l^\infty}| = \max\{l^m|\ P_0 \in l^m E(K)\}.$$

where the group $E(K[0])$ has no l^∞-torsion (as $l \in \mathcal{P}$, see Proposition 1.10.1 and Definition 3.1.3(f)).

We have the Hochschild-Serre spectral sequence

$$H^i(\mathrm{Gal}(K[0]/K), H^j(K[0], E_{l^n})) \Rightarrow H^{i+j}(K, E_{l^n}).$$

The short exact sequence of low degree terms of this spectral sequence is in part

$$0 \to H^1(\mathrm{Gal}(K[0]/K), E_{l^n}(K[0])) \to H^1(K, E_{l^n}) \to H^1(K[0], E_{l^n})^{\mathrm{Gal}(K[0]/K)}$$
$$\to H^2(\mathrm{Gal}(K[0]/K), E_{l^n}(K[0])).$$

The two extreme terms $H^1(\mathrm{Gal}(K[0]/K), E_{l^n}(K[0]))$ and $H^2(\mathrm{Gal}(K[0]/K), E_{l^n}(K[0]))$ are both zero as $E_{l^n}(K[0])$ is zero as already noted. Hence this short exact sequence provides the isomorphism

$$H^1(K, E_{l^n}) \cong H^1(K[0], E_{l^n})^{\mathrm{Gal}(K[0]/K)} \tag{5.1.3}$$

which is induced from the injection $E(K) \to E(K[0])$. The short exact sequence of sheaves for the étale topology on Spec K

$$0 \longrightarrow E_{l^n} \longrightarrow E \xrightarrow{l^n} E \longrightarrow 0$$

then provides the commutative diagram of cohomology groups

$$\begin{array}{ccccccccccc}
0 & \longrightarrow & E_{l^n}(K) & \longrightarrow & E(K) & \xrightarrow{l^n} & E(K) & \longrightarrow & H^1(K, E_{l^n}) & \ldots \\
 & & \downarrow & & \downarrow & & \downarrow & & \downarrow & \\
0 & \longrightarrow & E_{l^n}(K[0]) & \longrightarrow & E(K[0]) & \xrightarrow{l^n} & E(K[0]) & \longrightarrow & H^1(K[0], E_{l^n}) & \ldots
\end{array}$$

The isomorphism $H^1(K, E_{l^n}) \cong H^1(K[0], E_{l^n})^{\mathrm{Gal}(K[0]/K)}$ of (5.1.3) together with this commutative diagram then shows that the homomorphism

$$E(K)/l^n E(K) \to E(K[0])/l^n E(K[0])$$

is injective. Hence these two numbers l^{M_0} and $|(E(K)/\mathbb{Z}P_0)_{l^\infty}|$ are equal.

We obtain that the order of $\gamma_{M_0+m}(0)$ is equal to l^m from Proposition 3.4.14(i). □

5.1.4. Lemma. *Assume that the prime number $l \in \mathcal{P}$ is coprime to the order of* $\mathrm{Pic}(A)$. *Suppose that M_r is finite for some integer $r \geq 0$. Then M_s is finite for all $s \geq r$ and*

$$M_r, M_{r+1}, M_{r+2} \dots$$

is a decreasing sequence of non-negative integers.

Proof of Lemma 5.1.4. Suppose that M_s is finite for some $s \geq 0$. Then there is a divisor

$$c \in \Lambda^s(M_s + 1)$$

which satisfies $l^{M_s} || P_c$ and $M_s < \alpha(c)$; hence the cohomology class $\gamma_{M_s+1}(c)$ is non-zero (by Proposition 3.4.14(i)). From Proposition 3.3.8, where we take $n = M_s + 1$, we obtain a prime divisor

$$z \in \Lambda^1(M_s + 1) \tag{5.1.5}$$

coprime to c such that the localization $\gamma_{M_s+1}(c)_y$ is non-zero where y is the unique prime of K lying over z (see remark 3.2.5(1)). Then by Proposition 4.5.1(ii), we have $P_c \notin l^{M_s+1}E(K_y)$; that is to say

$$\mathrm{ord}((P_c \pmod{l^{M_s+1}})_y) > 1. \tag{5.1.6}$$

It follows from Proposition 4.5.1(iv) that $\mathrm{ord}\ \gamma_n(c+z)_y > 1$, where $n = M_s + 1$, and in particular we have $\gamma_n(c+z) \neq 0$. Therefore by Proposition 3.4.14(i)), or the definition of $\gamma_n(c+z)$, we have $P_{c+z} \notin l^{M_s+1}E(K[c+z])$. It follows that we have

$$c + z \in \Lambda^{s+1}(M_s + 1)$$

and M_{s+1} is finite and we have $M_{s+1} \leq M_s$. As M_r is finite by hypothesis, we then have that M_s is finite and $M_s \geq M_{s+1}$ for all $s \geq r$ by induction. □

5.1.7. Lemma. *Suppose that M_{r-1} is finite where $r \geq 1$. Assume that the prime number $l \in \mathcal{P}$ is coprime to the order of* $\mathrm{Pic}(A)$. *Let $c \in \Lambda^r(M_{r-1})$ and put $\nu(c) = (-1)^r\epsilon$. Then we have*
(i) $\delta_{M_{r-1}}(c) \in Ш(E/K)_{l^\infty}^{-\nu(c)}$
(ii) $\gamma_{M_{r-1}}(c) \in \mathrm{Sel}_{l^\infty}(E/K)^{-\nu(c)}$;
(iii) *the order of $\delta_{M_{r-1}}(c)$ is at most $l^{M_{r-1}-M_r}$.*

Proof of Lemma 5.1.7. By the previous Lemma 5.1.4, we have that M_r is finite and $M_{r-1} \geq M_r$. The set $\Lambda^r(M_{r-1})$ is non-empty by corollary 3.3.9; therefore

there is an element $c \in \Lambda^r(M_{r-1})$. It follows from the definition of the M_i that $l^{M_r}|P_c$ and $l^{M_{r-1}}|P_{c-z}$ for all prime divisors z in the support of c. From Proposition 4.5.1(iii) and Lemma 4.2.2 we obtain that

$$\delta_{M_{r-1}}(c) \in \text{Ш}(E/K)^{-\nu(c)}$$

Hence $\gamma_{M_{r-1}}(c)$ belongs to the Selmer group $\text{Sel}_{l^\infty}(E/K)^{-\nu(c)}$ (see Lemma 4.2.2). The order of the element $\delta_{M_{r-1}}(c)$ is at most $l^{M_{r-1}-M_r}$ by Proposition 3.4.14(ii). □

5.1.8. Lemma. *Let $z \in \Lambda^1(n)$ be a prime divisor. Let y be the unique prime of K lying above z.*
(i) *The Tate pairing on $H^1(K_y, E_{l^n})$ of Theorem 2.1.4 induces a non-degenerate pairing, where $\delta = \pm 1$,*

$$({}_{l^n}E(K_y))^\delta \times H^1(K_y, E)^\delta_{l^n} \longrightarrow \mathbb{Z}/l^n\mathbb{Z}$$

of eigenspaces under the action of τ which are finite cyclic groups of order l^n.
(ii) *The image of the homomorphism $\chi_z : {}_{l^n}E(K_y) \to H^1(K_y, E_{l^n})$ is an isotropic subgroup for the alternating Tate pairing on $H^1(K_y, E_{l^n})$ and $\text{Im}(\chi_z)^\delta \cong \mathbb{Z}/l^n\mathbb{Z}$ for $\delta = \pm 1$.*

Proof of Lemma 5.1.8. (i) We have isomorphisms of $\text{Gal}(K/F)$-modules

$${}_{l^n}E(K_y) \cong E(K_y)_{l^n}, \quad H^1(K_y, E)_{l^n} \cong \text{Hom}(\mu_{l^n}(K_{l^n}), E(K_y)).$$

The first isomorphism here follows from Proposition 4.4.1. As $|\kappa(z)|+1 \equiv 0 \pmod{l^n}$ and $\kappa(y)^*$ has $|\kappa(z)|^2 - 1$ elements we have that $\mu_{l^n}(K_y)$ has l^n elements and is contained in the -1 eigenspace under τ of $\kappa(y)^*$. Hence we have for $\delta = \pm 1$

$${}_{l^n}E(K_y)^\delta \cong H^1(K_y, E)^{-\delta}_{l^n}$$

and these groups are cyclic of order l^n by Lemma 3.2.7(ii).
(ii) The map χ_z is defined in Proposition 4.4.2. We have by property (3) of χ_z of this Proposition 4.4.2 that the composite of χ_z with the surjective homomorphism $H^1(K_y, E_{l^n}) \to H^1(K_y, E)_{l^n}$ is an isomorphism. This implies that that we have for $\delta = \pm 1$, where $\text{Im}(\chi_z)$ denotes the image of χ_z in $H^1(K_y, E_{l^n})$,

$$\text{Im}(\chi_z)^\delta \cong H^1(K_y, E)^\delta_{l^n}.$$

Hence part (i) on Tate duality implies that for $\delta = \pm 1$

$$\text{Im}(\chi_z)^\delta \cong \mathbb{Z}/l^n\mathbb{Z}.$$

This evidently shows that $\text{Im}(\chi_z)^\delta$ is an isotropic subgroup of $H^1(K_y, E_{l^n})$ for the anti-symmetric Tate pairing. Since the cup product on $H^1(K_y, E_{l^n})$ is $\text{Gal}(K/F)$-equivariant it follows that $\text{Im}(\chi_z) \cong \text{Im}(\chi_z)^{+1} \oplus \text{Im}(\chi_z)^{-1}$ is an isotropic subgroup of $H^1(K_y, E_{l^n})$. □

5.1.9. Lemma. *Let $z \in \Lambda^1(n)$ be a prime divisor where $n \geq 1$. Let y be the unique prime of K lying above z. Let S be a finite set of prime divisors of $\Lambda^1(n)$ not containing z. Let $\delta = \pm 1$. Then there is a non-zero element $h \in H^1(K, E_{l^n})^\delta$ in the δ-eigenspace satisfying these two conditions:*

(a) *$h_x \in \partial_{l^n}(E(K_x))$ for any place x of K not lying over a place of $S \cup \{z\}$;*

(b) *$h_x \in \mathrm{Im}(\chi_w)$ for all $w \in S$ where x is the unique place of K over w and where $\chi_w : {}_{l^n}E(K_x) \to H^1(K_x, E_{l^n})$ is the homomorphism of Proposition 4.4.2.*

Proof of Lemma 5.1.9. The places of S remain inert in the field extension K/F; for any $u \in S$ denote by $u^\sharp$ the corresponding place of K over u.

For any place $v \in \Sigma_K$ of K we have the exact sequence

$$0 \longrightarrow {}_{l^n}E(K_v) \xrightarrow{\partial_{l^n}} H^1(K_v, E_{l^n}) \longrightarrow H^1(K_v, E)_{l^n} \longrightarrow 0 \tag{5.1.10}$$

where the extremities of this sequence are in duality by Theorem 2.1.6.

Let $\delta = \pm 1$. The elliptic curve E/F has good reduction at all places of $S \cup \{z\}$ (remark 3.2.5). Let U be the finite subset of Σ_F of places of F given by

$$U = S \cup \{z\} \cup \{\text{bad reduction places of } E/F \text{ in } \Sigma_F\}.$$

Let U_K be the finite set of all places of K over the places of U. For all $v \in U_K$ dividing an element u of $U \setminus \{z\}$ put

$$H_v = \mathrm{Im}(\chi_u)^\delta \text{ if } u \in S$$

$$H_v = \partial_{l^n}(E(K_v))^\delta \text{ if } u \in U \setminus (S \cup \{z\}).$$

Then we have

$$|H_v|^2 = |H^1(K_v, E_{l^n})^\delta| \text{ for all } v \text{ dividing a place } u \in U \setminus \{z\}.$$

For the case where $u \in S$ this follows from Lemma 5.1.8(ii). For the case where $u \in U \setminus (S \cup \{z\})$, that is to say a place of bad reduction, this follows from the exact sequence (5.1.10) and that the extremities of this sequence are in duality.

From Tate local and global duality, there is a self dual exact sequence (see [9, Chap. 1, Theorem 4.10, p.70])

$$H^1(K_U/K, E_{l^n}) \to \bigoplus_{v \in U_K} H^1(K_v, E_{l^n}) \to H^1(K_U/K, E_{l^n})^*$$

where N^*, for a $\mathbb{Z}/l^n\mathbb{Z}$-module N, denotes $\mathrm{Hom}(N, \mathbb{Z}/l^n\mathbb{Z})$, and where K_U is the maximal extension of K unramified outside U_K. Hence the image I of $H^1(K_U/K, E_{l^n})$ in $\bigoplus_{v \in U_K} H^1(K_v, E_{l^n})$ is a maximal isotropic subgroup, for the Tate pairing, of the group

$$\bigoplus_{v \in U_K} H^1(K_v, E_{l^n}).$$

Since at the place y lying over z we have $H^1(K_y, E_{l^n})^\delta \neq 0$ by Lemma 5.1.8(i), the subgroup I^δ is of order strictly larger than that of

$$\bigoplus_{u\in S} \frac{H^1(K_{u^\sharp}, E_{l^n})^\delta}{H_{u^\sharp}}.$$

Hence the natural homomorphism

$$H^1(K_U/K, E_{l^n})^\delta \to \bigoplus_{u\in S} \frac{H^1(K_{u^\sharp}, E_{l^n})^\delta}{H_{u^\sharp}}$$

has non-zero kernel. Therefore we may select a non-zero element $h \in H^1(K_U/K, E_{l^n})^\delta$ in this kernel. Then h satisfies the condition (b), that is to say we have $h_{u^\sharp} \in H_{u^\sharp}$ for all $u \in S$. Furthermore, h satisfies the condition (a) by the selection of H_v if $v \in \Sigma_K$ is a bad reduction place of E and because we have $H^1(K_v^{\mathrm{un}}/K_v, E_{l^n}) = \partial_{l^n}(E(K_v))$ if $v \in \Sigma_K$ is a good reduction place of E, where K_v^{un} is the maximal unramified separable extension of K_v, □

5.1.10. Remark. The Lemma 5.1.9 is a technical result required for the proof of Proposition 5.2.1 in the next section.

5.2. A class $\gamma_n(c)$ in the Selmer group

The notation and hypotheses (5.1.1) of section 5.1 hold also for this section.

5.2.1. Proposition. *Let $r \geq 1$ be an integer and put $\nu(r) = (-1)^r\epsilon$. Let G be a subgroup of the Selmer eigenspace $\mathrm{Sel}_{l^\infty}(E/K)^{-\nu(r)}$ such that*

$$\mathrm{rank}(G) \leq r.$$

Assume that M_{r-1} is finite and that $l \in \mathcal{P}$ is coprime to the order of $\mathrm{Pic}(A)$. Then there is a cohomology class $\gamma_{M_{r-1}}(c)$ for some divisor $c \in \Lambda^r(M_{r-1})$ such that:
(i) *$\gamma_{M_{r-1}}(c)$ belongs to the Selmer eigenspace $\mathrm{Sel}_{l^\infty}(E/K)^{-\nu(r)}$;*
(ii) *$\gamma_{M_{r-1}}(c)$ has order $l^{M_{r-1}-M_r}$;*
(iii) $\mathbb{Z}\gamma_{M_{r-1}}(c) \cap G = \{0\}$.

Proof of Proposition 5.2.1. The group $\mathrm{Sel}_{l^\infty}(E/K)$ is an l^∞-torsion group, that is to say every element is annihilated by a power of l. Hence the subgroup G of finite rank is therefore finite.

Let $\exp(G)$ be the exponent of G, that is to say the largest order of an element of the abelian l-group G. Select an integer

$$m \geq 1$$

such that

$$l^m \geq \max\{\exp(G), l^{M_{r-1}}\}$$

and put

$$L = K(E_{l^m}).$$

Put

$$t = \mathrm{rank}(G)$$

where

$$\mathrm{rank}(G) = \dim_{\mathbb{Z}/l\mathbb{Z}} G/lG$$

and where $t \leq r$ by hypothesis. Note that G is a subgroup of $\mathrm{Hom}(\mathrm{Gal}(L^{\mathrm{sep}}/L), E_{l^m}(L))$, as $l^m \geq \exp(G)$ (see Lemma 3.3.2), and the elements of $\Lambda^1(m)$ are unramified in L/F. As in §1.2, for any divisor c on F, $\mathrm{Supp}(c)$ denotes the set of distinct prime divisors in the support of c.

For any divisor c in $\Lambda^r(1)$, put

$$\Xi(c) = \mathrm{Supp}(c) \cap \Lambda^1(m)$$

that is to say $\Xi(c)$ is the set of prime divisors in the support of c which belong to $\Lambda^1(m)$; in particular, $\Xi(c)$ depends on m. For any prime divisor z of F in $\Lambda^1(1)$, select a place $z^\times$ of $L = K(E_{l^m})$ lying over z. Let $\Gamma(c) \subseteq \widehat{G}$ be the subgroup of characters of the abelian group G generated by the set of characters

$$\{\phi_{\mathrm{Frob}(z^\times)} \quad | \quad z \in \Xi(c)\}$$

(as in Proposition 3.3.3 and (3.3.4), applied to the finite group G, and Proposition 3.3.7). Put

$$s(c) = \dim_{\mathbb{Z}/l\mathbb{Z}} \frac{\Gamma(c) + l\,\widehat{G}}{l\,\widehat{G}}.$$

That is to say, $s(c)$ is the dimension of the image of $\Gamma(c)$ in the vector space $\widehat{G}/l\,\widehat{G}$ of dimension t; the non-negative integer $s(c)$ is then at most equal to t. Put

$$n(c) = |\Xi(c)|$$

that is to say $n(c)$ is the cardinality of $\Xi(c)$ and which is a non-negative integer at most equal to r, the number of prime divisors in the support of c.

Define the *defect* $\Delta(c)$ of a divisor $c \in \Lambda^r(1)$ on F to be

$$\Delta(c) = \max(t - s(c), r - n(c)).$$

Then we have

$$0 \leq \Delta(c) \leq r$$

and we have

$$\Delta(c) = 0 \text{ if and only if } \Gamma(c) = \widehat{G} \text{ and } c \in \Lambda^r(m).$$

This equivalence holds because $s(c) = t$ if and only if $\Gamma(c) = \widehat{G}$ by Nakayama'a Lemma.

We have $M_{r-1} \geq M_r$ by Lemma 5.1.4 and in particular M_r is finite as M_{r-1} is assumed to be finite. We may then select a divisor

$$d \in \Lambda^r(M_r + 1) \tag{5.2.2}$$

such that

$$l^{M_r} || P_d.$$

Then the cohomology class $\gamma_{M_r+1}(d)$ is defined and belongs to $H^1(K, E_{l^{M_r+1}})^{-\nu(r)}$ and has exact order l by Lemma 4.2.2 and Proposition 3.4.14(i).

Suppose that $M_r = M_{r-1}$. Then $\gamma_{M_{r-1}}(d)$ is equal to zero and evidently belongs to the Selmer eigenspace $\mathrm{Sel}_{l^\infty}(E/K)^{-\nu(r)}$ which proves the lemma in this trivial case where $M_r = M_{r-1}$.

We may suppose for the rest of the proof of this Proposition 5.2.1 that $M_{r-1} > M_r$.

Assume that the defect of the divisor d already selected in (5.2.2) satisfies

$$\Delta(d) > 0.$$

That is to say either (where $s(d) < t$) the image of $\Gamma(d)$ is a proper subspace of $\widehat{G}\,/l\,\widehat{G}$ or (where $n(d) < r$) that $d \notin \Lambda^r(m)$. These two possibilities $s(d) < t$ and $n(d) < r$ for the divisor d are not mutually exclusive.

We select a character $\psi \in \widehat{G}$ and a prime divisor $z_0 \in \mathrm{Supp}(d)$ by the following recipe.

Selection of the character ψ

If $s(d) < t$ select a character $\psi \in \widehat{G}$ such that

$$\psi(\gamma_{M_r+1}(d)) \neq 0, \quad \text{if } \gamma_{M_r+1}(d) \in G \tag{5.2.3}$$

and

$$\psi \in \widehat{G} \setminus (\Gamma(d) + l\,\widehat{G}) \tag{5.2.4}$$

where the condition (5.2.3) is vacuous if $\gamma_{M_r+1}(d) \notin G$. A character $\psi \in \widehat{G}$ exists which satisfies the two conditions (5.2.3) and (5.2.4) because a finite group cannot be the union of two proper subgroups.

If $s(d) = t$ and $n(d) < r$ select any character $\psi \in \widehat{G}$ such that

$$\psi(\gamma_{M_r+1}(d)) \neq 0, \quad \text{if } \gamma_{M_r+1}(d) \in G \tag{5.2.5}$$

where the condition (5.2.5) is vacuous if $\gamma_{M_r+1}(d) \notin G$.

Selection of the divisor $z_0 \in \mathrm{Supp}(d)$

If $s(d) < t$ select a prime divisor $z_0 \in \mathrm{Supp}(d)$ such that

$$\{\phi_{\mathrm{Frob}(z^\times)} \quad | \quad z \in \Xi(d - z_0)\}$$

is a generating set for $\Gamma(d)$ modulo $l\widehat{G}$. This choice is possible because $t \leq r$ and $\mathrm{Supp}(d)$ has r distinct elements.

If $s(d) = t$ and $n(d) < r$ select a prime divisor $z_0 \in \mathrm{Supp}(d)$ such that $z_0 \notin \Lambda^1(m)$.

We have now defined the pair ψ, z_0 for the divisor d where the defect $\Delta(d) > 0$. By Lemma 5.1.9 we may select a cohomology class h in the $\nu(r)$-eigenspace

$$h \in H^1(K, E_l)^{\nu(r)}$$

of the group scheme E_l such that

(5.2.6) $h \neq 0$;
(5.2.7) $h_v \in \partial_l(E(K_v))$ for all places v of K coprime to $\mathrm{Supp}(d)$;
(5.2.8) $h_y \in \mathrm{Im}(\chi_z)$ for all $z \in \mathrm{Supp}(d - z_0)$ where y is the prime of K lying over z and χ_z is the homomorphism of §4.4.

Note that $H^1(K, E_l)^{\nu(r)}$ is a subgroup of $H^1(K, E_{l^m})$ by Lemma 3.1.5.

Since h is in a different eigenspace from G and $\gamma_{M_r+1}(d)$, which both belong to the $-\nu(r)$-eigenspace, we have

$$(G + \mathbb{Z}\gamma_{M_r+1}(d)) \cap \mathbb{Z}h = 0$$

where $G + \mathbb{Z}\gamma_{M_r+1}(d)$ is the subgroup of $H^1(K, E_{l^m})^{-\nu(r)}$ generated by G and $\gamma_{M_r+1}(d)$ and where $\mathbb{Z}h$ is the subgroup of $H^1(K, E_{l^m})^{\nu(r)}$ generated by h.

Let D be the subgroup of $H^1(K, E_{l^m})$ generated by G, $\gamma_{M_r+1}(d)$ and h

$$D = G + \mathbb{Z}\gamma_{M_r+1}(d) + \mathbb{Z}h.$$

Then D is a finite subgroup of $H^1(K, E_{l^m})$ by Lemma 3.1.5. As D is isomorphic to the direct product of $G + \mathbb{Z}\gamma_{M_r+1}(d)$ and $\mathbb{Z}h$, we can select a homomorphism

$$\chi : D \to \mathbb{Q}/\mathbb{Z}$$

such that (by (5.2.3), (5.2.5), and (5.2.6))

(5.2.9) $\chi|_G = \psi$;
(5.2.10) $\chi(\gamma_{M_r+1}(d)) \neq 0$;
(5.2.11) $\chi(h) \neq 0$.

By Proposition 3.3.7 applied to the finite group D, there is a prime divisor

$$z_1 \in \Lambda^1(m) \tag{5.2.12}$$

distinct from the elements of $\mathrm{Supp}(d)$ such that, in the notation of (3.3.4),

$$\chi = \phi_{\mathrm{Frob}(z_1^{\times})} \tag{5.2.13}$$

where $z_1^{\times}$ is a prime divisor of $L = K(E_{l^m})$ above z_1. We then obtain the cohomology class $\gamma_{M_r+1}(d+z_1)$ associated to the divisor $d+z_1$.

For all places v of K, denote the alternating cup product induced by the Weil pairing of h with $\gamma_{M_r+1}(d+z_1)$ localized at v by (see Theorem 2.1.4)

$$< \gamma_{M_r+1}(d+z_1)_v, h_v >_v$$

which is an element of $\mathbb{Q}/\mathbb{Z}$. The sum of local pairings over all places of K

$$\sum_{\text{all places } v \text{ of } K} < \gamma_{M_r+1}(d+z_1)_v, h_v >_v = 0 \tag{5.2.14}$$

is zero by Proposition 2.1.8.

If v does not divide any element of $\mathrm{Supp}(d+z_1)$ then

$$\gamma_{M_r+1}(d+z_1)_v \in \partial_{l^{M_r+1}}(E(K_v))$$

by Proposition 4.5.1(i) and

$$h_v \in \partial_l(E(K_v))$$

by (5.2.7), that is to say both localized elements $\gamma_{M_r+1}(d+z_1)_v$, h_v are in the image of the map

$$\partial_{l^m} : E(K_v) \to H^1(K_v, E_{l^m}).$$

But the image of this map ∂_{l^m} in $H^1(K_v, E_{l^m})$ is an isotropic subgroup for the alternating pairing $<,>_v$ (see [1, p. 403, Theorem 7.15.6] or Theorem 2.1.6(i)). Therefore we have

$$< \gamma_{M_r+1}(d+z_1)_v, h_v >_v = 0 \text{ for all } v \text{ coprime to } \mathrm{Supp}(d+z_1).$$

If y is a place of K which divides an element z of $\mathrm{Supp}(d-z_0)$, then we have $h_y \in \mathrm{Im}(\chi_z)$ by (5.2.8) and

$$\gamma_{M_r+1}(d+z_1)_y \in \mathrm{Im}(\chi_z)$$

by property (4) of Proposition 4.4.2, which is the main property of the map χ_z. As $\mathrm{Im}(\chi_z)$ is isotropic for the cup product $<,>_v$ by Lemma 5.1.8(ii), we have

$$< \gamma_{M_r+1}(d+z_1)_y, h_y >_y = 0 \text{ for all } y \text{ dividing an element of } \mathrm{Supp}(d-z_0).$$

Therefore the only possible non-zero terms in the sum $\sum_v < \gamma_{M_r+1}(d+z_1)_v, h_v >_v$ of (5.2.14) are for the places of K lying over the places z_0 and z_1. For these places $z_i \in \Lambda^1$ of F, for $i = 0, 1$, denote by y_i the corresponding place of K lying over the place z_i which remains inert in K/F.

From (5.2.10) and (3.3.5) and that $\chi = \phi_{\mathrm{Frob}(z_1^\times)}$ by (5.2.13), we have that this localization $\gamma_{M_r+1}(d)_{y_1}$ at y_1 is non-zero; hence by Proposition 4.5.1(iv) the localization

$$\delta_{M_r+1}(d+z_1)_{y_1} \in H^1(K_{y_1}, E)^{\nu(r)}_{l^{M_r+1}}$$

is non-zero. Furthermore we have

$$h_{y_1} \in \partial_l(E(K_{y_1}))^{\nu(r)}$$

by (5.2.7) and h_{y_1} is non-zero by (5.2.11) and (5.2.13). Hence we have by Lemma 5.1.8(i)

$$< \gamma_{M_r+1}(d+z_1)_{y_1}, h_{y_1} >_{y_1} \neq 0.$$

Since the sum $\sum_v < \gamma_{M_r+1}(d+z_1)_v, h_v >_v$ of (5.2.14) is zero, this implies that the local term at y_0

$$< \gamma_{M_r+1}(d+z_1)_{y_0}, h_{y_0} >_{y_0}$$

is non-zero. Hence we have $\gamma_{M_r+1}(d+z_1)_{y_0} \neq 0$ and so by Proposition 4.5.1(iv)

$$P_{d+z_1-z_0} \notin l^{M_r+1}E(K_{y_0})$$

hence we obtain, where $z_1 \in \Lambda^1(m)$,

$$l^{M_r} || P_{d+z_1-z_0} \tag{5.2.15}$$

because we have $l^{M_r} | P_{d+z_1-z_0}$ and $d+z_1-z_0 \in \Lambda^r(M_r+1)$ as well as $d \in \Lambda^r(M_r+1)$ and $z_1 \in \Lambda^1(m)$.

Put

$$c_1 = d + z_1 - z_0 \in \Lambda^r(M_r+1).$$

On the one hand, we have that if $s(d) < t$ then the group $\Gamma(c_1)$ is generated by $\Gamma(d)$ and ψ

$$\Gamma(c_1) = \Gamma(d) + \mathbb{Z}\psi \tag{5.2.16}$$

because $\chi = \phi_{\mathrm{Frob}(z_1^\times)}$ by (5.2.13) and $\chi|_G = \psi$ by (5.2.9) in this case where $s(d) < t$. The condition that $z_1 \in \Lambda(m)$ is satisfied by the choice of z_1 in (5.2.12). We then have from (5.2.16) if $s(d) < t$, as $\psi \notin \Gamma(d) + l\widehat{G}$ by (5.2.4),

$$s(c_1) = \dim_{\mathbb{Z}/l\mathbb{Z}} \frac{\Gamma(c_1) + l\widehat{G}}{l\widehat{G}} = s(d) + 1.$$

On the other hand, if $s(d) = t$ and $n(d) < r$ then $n(c_1) = n(d) + 1$ as $z_1 \in \Lambda^1(m)$ by (5.2.12) and $z_0 \notin \Lambda^1(m)$ by the selection of $z_0 \in \mathrm{Supp}(d)$. Hence we have in both cases that the defect of c_1 is given by

$$\Delta(c_1) = \Delta(d) - 1.$$

We may by this method construct by induction a sequence of divisors $c_1, c_2, c_3 \ldots$ in $\Lambda^r(M_r + 1)$ such that their defects are strictly decreasing

$$\Delta(d) > \Delta(c_1) > \Delta(c_2) > \ldots$$

and where, as in (5.2.15),

$$l^{M_r} || P_{c_i} \text{ for all } i. \tag{5.2.17}$$

This sequence must terminate in a divisor c with zero defect $\Delta(c) = 0$, that is to say $\Gamma(c) = \widehat{G}$, by Nakayama's Lemma, and $c \in \Lambda^r(m)$ and where

$$l^{M_r} || P_c. \tag{5.2.18}$$

The cohomology class $\gamma_m(c)$ is therefore defined and as $\Gamma(c) = \widehat{G}$ we have, where $z^\times$ is a place of $K(E_{l^m})$ over z,

$$\{g \in G |\ \phi_{\mathrm{Frob}(z^\times)}(g) = 0 \text{ for all } z \in \mathrm{Supp}(c)\} = 0. \tag{5.2.19}$$

We have from Lemma 5.1.7(ii) that

$$\gamma_{M_{r-1}}(c) \in \mathrm{Sel}_{l^\infty}(E/K)^{-\nu(r)}. \tag{5.2.20).}$$

We have by the definition of M_{r-1} that $\gamma_{M_{r-1}}(c - z) = 0$ for any prime divisor z in the support of c as $c - z$ has $r - 1$ elements in its support. But by Proposition 4.5.1(iv), we have for any $z \in \mathrm{Supp}(c)$ where y is the place of K over z,

$$\mathrm{ord}\ \gamma_{M_{r-1}}(c)_y = \mathrm{ord}\ \gamma_{M_{r-1}}(c - z)_y.$$

Hence we obtain for any $z \in \mathrm{Supp}(c)$ that $\gamma_{M_{r-1}}(c)_y = 0$ where y is the prime of K over z. We then obtain from (5.2.19), as $g_y = \phi_{\mathrm{Frob}(z^\times)}(g)$ where y is the place of K above z, that

$$G \cap \mathbb{Z}\gamma_{M_{r-1}}(c) = 0.$$

Since $l^{M_r} || P_c$, by (5.2.18), we have that $\gamma_m(c)$ has order $l^{m - M_r}$ for all $m \geq M_r$ by Proposition 3.4.14(i). Hence $\gamma_{M_{r-1}}(c)$ has order $l^{M_{r-1} - M_r}$ and belongs to the Selmer group $\mathrm{Sel}_{l^\infty}(E/K)^{-\nu(r)}$, by (5.2.20), which proves the proposition. □

5.3. Proof of Theorem 4.1.15

(5.3.1) The notation (5.1.1) of section §5.1 holds also for this section.

5.3.2. Lemma. *Let A be a finite abelian p-group where p is a prime number and with invariants $I_1 \geq I_2 \geq \ldots \geq I_r$. Let B be a subgroup of A with invariants $I_1 \geq I_2 \geq \ldots \geq I_s$ where $s \leq r$. Then there is a subgroup C of A such that*

$$A = B \oplus C$$

that is to say, A is the direct sum of the subgroups B, C. The invariants of C are $I_{s+1} \geq I_{s+1} \geq \ldots \geq I_r$.

[The proof of this result follows from the structure theorem of finite abelian groups and is omitted.] □

For the proof of Theorem 4.1.15, as P_0 has infinite order in $E(K)$, the Tate-Shafarevich group $Ш(E/K)$ is finite and P_0 generates a subgroup of finite index in $E(K)$ by [1, Chapter 7, Theorem 7.6.5] which is one of the principal results of this monograph [1].

The image of P_0 in ${}_{l^m}E(K)$ belongs to the $-\epsilon$-eigenspace of ${}_{l^m}E(K)$ for all m (by [1, Chapter 7, Lemma 7.14.11] or Lemma 4.2.2 above). It follows from the decomposition into eigenspaces

$${}_{l^m}E(K) \cong ({}_{l^m}E(K))^{\epsilon} \oplus ({}_{l^m}E(K))^{-\epsilon}$$

that the ϵ-eigenspace $({}_{l^m}E(K))^{\epsilon}$ is a finite abelian group of order bounded independently of m and hence $E(K)^{\epsilon}$ is a finite abelian group. As $E(K)$ has no l-torsion (as $l \in \mathcal{P}$ see Proposition 1.10.1 and Definition 3.1.3(f)) it follows that

$$({}_{l^m}E(K))^{\epsilon} = 0 \quad \text{for all } m \tag{5.3.3}$$

and that

$$({}_{l^m}E(K))^{-\epsilon} \cong \mathbb{Z}/l^m\mathbb{Z} \ \text{ for all } m \tag{5.3.4}$$

which proves the isomorphisms (4.1.18).

The l^m-Selmer group $\mathrm{Sel}_{l^m}(E/K)^{\pm}$ belongs to an exact sequence of eigenspaces

$$0 \longrightarrow ({}_{l^m}E(K))^{\pm} \longrightarrow \mathrm{Sel}_{l^m}(E/K)^{\pm} \longrightarrow Ш(E/K)^{\pm}_{l^m} \longrightarrow 0. \tag{5.3.5}$$

Hence this exact sequence induces an isomorphism of ϵ-eigenspaces

$$\mathrm{Sel}_{l^\infty}(E/K)^{\epsilon} \cong Ш(E/K)^{\epsilon}_{l^\infty}. \tag{5.3.6}$$

The largest invariant of the $-\epsilon$-eigenspace $\mathrm{Sel}_{l^m}(E/K)^{-\epsilon}$ is at most equal to m. But this group $\mathrm{Sel}_{l^m}(E/K)^{-\epsilon}$ contains the subgroup $({}_{l^m}E(K))^{-\epsilon} \cong \mathbb{Z}/l^m\mathbb{Z}$

with invariant m. By Lemma 5.3.2 applied to the eigenspace $\mathrm{Sel}_{l^m}(E/K)^{-\epsilon}$ and the subgroup $({}_{l^m}E(K))^{-\epsilon}$, we have that the $-\epsilon$-eigencomponent of the exact sequence (5.3.5) splits. Hence we obtain an isomorphism of $-\epsilon$-eigenspaces

$$\mathrm{Sel}_{l^m}(E/K)^{-\epsilon} \cong {}_{l^m}E(K) \oplus \text{Ш}(E/K)_{l^m}^{-\epsilon} \quad \text{for all } m. \tag{5.3.7}$$

The isomorphisms (5.3.6) and (5.3.7) prove the theorem. □

5.4. Proof of Theorems 4.1.9 and 4.1.13

(5.4.1) The principle of the proof is to construct inductively divisors $c_k \in \Lambda^k$, $k = 1, 2 \ldots$, such that the cohomology classes $\delta_{M_{k-1}}(c_k)$ in the Tate-Shafarevich group $\text{Ш}(E/K)_{l^\infty}$ form a basis of a maximal isotropic subgroup with respect to the Cassels pairing and where $\delta_{M_{k-1}}(c_k)$ has order l^{N_k} for all k. The inductive step is provided by the next Lemma 5.4.3.

(5.4.2) The notation and hypotheses (5.1.1) of the previous section holds also for this section. We further denote by, where $l \in \mathcal{P}$,

Σ_K the set of all places of the global field K;
$[,]_v : {}_{l^m}E(K_v) \times H^1(K_v, E)_{l^m} \to \mathbb{Z}/l^m\mathbb{Z}$ the non-degenerate local pairing at the place v of K induced by the cup product, as in Theorem 2.1.6.

If $z_i \in \Lambda^1(1)$ is a prime divisor of F indexed by an integer i, denote by y_i the prime divisor of K lying above the place z_i of F where this place z_i is inert in the field extension K/F.

5.4.3. Lemma. *Assume that $l \in \mathcal{P}$ is coprime to the order of $\mathrm{Pic}(A)$. Let $s \geq 1$ be a positive integer and let $r, t \geq 0$ be non-negative integers. Let S be a subgroup of $\mathrm{Sel}_{l^s}(E/K)$. Let $e \in \mathrm{Sel}_{l^s}(E/K)^{-\nu(r+1)}$, $\gamma_s(c) \in \mathrm{Sel}_{l^s}(E/K)^{-\nu(r)}$, where $c \in \Lambda^r(s+t)$, be elements of the Selmer group where*

$$S \cap (e, \gamma_s(c)) = 0$$

and where $(e, \gamma_s(c))$ is the subgroup of the Selmer group generated by $e, \gamma_s(c)$. Suppose also that $e, \gamma_s(c)$ both have order l^n where $n \leq s$. Then there are infinitely many prime divisors $z \in \Lambda^1(s+t)$ coprime to $\mathrm{Supp}(c)$ such that if y is the place of K over z then we have
(1) $S_y = 0$;
(2) the value in $\mathbb{Z}/l^n\mathbb{Z}$ of the local pairing at y with the class $\delta_s(c+z)$

$$[\delta_s(c+z)_y, w_y]_y$$

has order l^n where $w_y \in {}_{l^n}E(K_y)$ is a point such that $e_y = \partial_{l^n}(w_y)$.

Proof of Lemma 5.4.3. Note that from (3.1.10), we have that $\mathrm{Sel}_{l^n}(E/K)$ is the subgroup of $\mathrm{Sel}_{l^s}(E/K)$ annihilated by l^n and in particular $\mathrm{Sel}_{l^n}(E/K)$ contains e and $\gamma_s(c)$.

Let T be the subgroup of the Selmer group $\mathrm{Sel}_{l^s}(E/K)$ generated by S, e and $\gamma_s(c)$. Then we have an isomorphism

$$T \cong S \oplus (e, \gamma_s(c)).$$

For a fixed non-zero element $x \in T$, the set of characters

$$\chi : T \to \mathbb{Z}/l^s\mathbb{Z}$$

such that

$$\mathrm{ord}(\chi(x)) < \mathrm{ord}(x)$$

is a proper subgroup of $\widehat{T}$. This follows as the subgroups of $\mathbb{Z}/l^s\mathbb{Z}$ are linearly ordered $\mathbb{Z}/l^s\mathbb{Z} \supseteq l\mathbb{Z}/l^s\mathbb{Z} \supseteq l^2\mathbb{Z}/l^s\mathbb{Z} \ldots$. As a group cannot be the union of two proper subgroups, there is then a character $\chi : T \to \mathbb{Z}/l^s\mathbb{Z}$ such that

$$\mathrm{ord}(\chi(e)) = \mathrm{ord}(e)$$

$$\mathrm{ord}(\chi(\gamma_s(c)) = \mathrm{ord}(\gamma_s(c))$$

$$\chi(S) = 0.$$

By Proposition 3.3.8 applied to the subgroup T and the character χ, we may select a prime divisor $z \in \Lambda^1(s+t)$ satisfying, where y is the place of K lying over z,

$$\mathrm{ord}(e_y) = \mathrm{ord}(e)$$

$$\mathrm{ord}(\gamma_s(c)_y) = \mathrm{ord}(\gamma_s(c))$$

$$S_y = 0$$

where the subscript y denotes the restriction at y of elements of the Selmer group $\mathrm{Sel}_{l^s}(E/K)$. In particular, condition (1) of the lemma is satisfied for this z.

The class $\delta_s(c+z)$, associated to the divisor $c+z \in \Lambda^{r+1}(s+t)$, then belongs to $H^1(K,E)_{l^s}^{-\nu(r+1)}$ and e belongs to $H^1(K, E_{l^s})^{-\nu(r+1)}$.

We may select a point $w_y \in ({}_{l^n}E(K_y))^{-\nu(r+1)}$ such that $e_y = \partial_{l^n}(w_y)$ where $e \in \mathrm{Sel}_{l^n}(E/K)$ as already noted. Then w_y has order l^n in ${}_{l^n}E(K_y)$ as e_y has order l^n.

Furthermore because

$$\mathrm{ord}(\gamma_s(c)_y) = \mathrm{ord}(\gamma_s(c))$$

and by Proposition 4.5.1(iv) we must have that

$$\mathrm{ord}(\delta_s(c+z)_y) = \mathrm{ord}(\gamma_s(c)) = l^n.$$

The class $\delta_s(c+z)_y$, of order l^n, belongs to the subgroup $H^1(K_y,E)_{l^n}^{-\nu(r+1)}$ of $H^1(K_y,E)_{l^s}^{-\nu(r+1)}$ and w_y, of order l^n, belongs to $({}_{l^n}E(K_y))^{-\nu(r+1)}$. In particular, $\delta_s(c+z)$ and w_y both belong to the $-\nu(r+1)$-eigenspaces of their respective spaces. Hence the local term

$$[\delta_s(c+z)_y, w_y]_y$$

has order l^n. This follows from the non-degeneracy of the local pairing: by Lemma 5.1.8(i), if $z \in \Lambda^1(n)$ and $y \in \Sigma_K$ is the place of K over z then the two elements

$$f \in {}_{l^n}E(K_y),\ d \in H^1(K_y,E)_{l^n}$$

give via the Tate pairing an element

$$[f,d]_y$$

which is non-zero if they are in the same τ-eigenspace and the product of their orders is $\geq l^n$. Therefore condition (2) is satisfied by z which proves the lemma.□

We now prove simultaneously the two Theorems 4.1.9 and 4.1.13.

As P_0 has infinite order in $E(K)$, the Tate-Shafarevich group $Ш(E/K)$ is finite and P_0 generates a subgroup of finite index in $E(K)$ by [1, Chapter 7, Theorem 7.6.5].

By definition of the invariants N_i of the finite abelian group $Ш(E/K)_{l^\infty}$ in §4.1.5, there is a maximal isotropic subgroup D of $Ш(E/K)_{l^\infty}$, with respect to the non-degenerate anti-symmetric Cassels pairing, where

$$D = \prod_i D_i$$

and each D_i is a cyclic group of order l^{N_i} and where

$$D^\epsilon = \prod_{i \text{ odd}} D_i$$

and

$$D^{-\epsilon} = \prod_{i \text{ even}} D_i.$$

From Theorem 4.1.15, we have the decomposition of eigenspaces

$$\text{Sel}_{l^m}(E/K)^\pm \cong ({}_{l^m}E(K))^\pm \oplus Ш(E/K)^\pm_{l^m} \quad \text{for all } m \geq 0 \tag{5.4.4}$$

where the projection onto the second factor is given by the natural surjection

$$\pi_m : \text{Sel}_{l^m}(E/K) \longrightarrow Ш(E/K)_{l^m}.$$

Let m be an integer such that l^m is greater than or equal to the exponent of the finite group $Ш(E/K)_{l^\infty}$, that is to say $m \geq \max_i N_i$. For each integer i, let $d_i \in Ш(E/K)_{l^\infty}^{-\nu(i)}$ be a generator of D_i and let e_i such that

$$e_i \in \mathrm{Sel}_{l^m}(E/K)^{-\nu(i)}, \text{ where } \pi_m(e_i) = d_i, \ \mathrm{ord}(e_i) = l^{N_i}, \tag{5.4.5}$$

be the lifting of d_i to the Selmer eigenspace $\mathrm{Sel}_{l^m}(E/K)^{-\nu(i)}$ via the decomposition (5.4.4) such that e_i has zero component in the subgroup ${}_{l^m}E(K)$; in particular, we take e_i to have order equal to l^{N_i} for all i and to belong to the $-\nu(i)$-eigenspace as d_i has order l^{N_i}. For each valuation v of K and each i, select $w_{i,v} \in {}_{l^{N_i}}E(K_v)$ such that the localization $e_{i,v}$ of e_i at v satisfies

$$e_{i,v} = \partial_{l^{N_i}}(w_{i,v}). \tag{5.4.6}$$

Here

$$\partial_{l^{N_i}} : {}_{l^{N_i}}E(K_v) \to H^1(K_v, E_{l^{N_i}}) \tag{5.4.7}$$

denotes the connecting homomorphism associated to the morphism $l^{N_i} : E \to E$ of multiplication by l^{N_i}.

The cohomology class $\gamma_{M_0+N_1}(0)$ belongs to the Selmer group $\mathrm{Sel}_{l^m}(E/K)^{-\epsilon}$ and has order l^{N_1} as $l^{M_0}||P_0$, by Lemma 5.1.2 or Proposition 3.4.14(i),

$$\mathrm{ord}(\gamma_{M_0+N_1}(0)) = l^{N_1}.$$

The element $e_1 \in \mathrm{Sel}_{l^m}(E/K)^{\epsilon}$ has the same order

$$\mathrm{ord}(e_1) = l^{N_1}.$$

Let S be the subgroup of the Selmer group $\mathrm{Sel}_{l^m}(E/K)$ generated by e_i for all $i \geq 2$. The element $\gamma_{M_0+N_1}(0)$ belongs to the component ${}_{l^m}E(K)$ in the decomposition (5.4.4) of $\mathrm{Sel}_{l^m}(E/K)$. Hence we have that the subgroup of $\mathrm{Sel}_{l^m}(E/K)$ generated by $S, e_1, \gamma_{M_0+N_1}(0)$ is direct sum

$$S \oplus \mathbb{Z}e_1 \oplus \mathbb{Z}\gamma_{M_0+N_1}(0).$$

We may now apply Lemma 5.4.3 to S and the elements $e_1, \gamma_{M_0+N_1}(0)$ where we take the parameters of the lemma to be

$$c = 0, r = 0, s = M_0 + N_1, t = 0, n = N_1. \tag{5.4.8}$$

There is according to the lemma a prime divisor $z_1 \in \Lambda^1(M_0 + N_1)$ which satisfies the following 2 conditions, where $y_1 \in \Sigma_K$ is the prime divisor of K lying over z_1 and where the subscript y_1 denotes localization at y_1, and where the point w_{1,y_1} in $(E(K_{y_1})/l^{N_1}E(K_{y_1}))^{\epsilon}$ is such that $\partial_{l^{N_1}}(w_{1,y_1}) = e_{1,y_1}$:-

$$\mathrm{ord}[\delta_{M_0+N_1}(z_1)_{y_1}, w_{1,y_1}]_{y_1} = l^{N_1} \tag{5.4.9}$$

and

$$S_{y_1} = 0. \tag{5.4.10}$$

Here in (5.4.9), $\delta_{M_0+N_1}(z_1)$ is the cohomology class in $H^1(K,E)^{\epsilon}_{l^{M_0+N_1}}$ associated to z_1.

Let

$$\delta_{M_0}(z_1) \in \text{Ш}(E/K)^{\epsilon}_{l^\infty}$$

be the cohomology class associated to this prime divisor $z_1 \in \Lambda^1(M_0+N_1)$; that $\delta_{M_0}(z_1)$ belongs to the Tate-Shafarevich group $\text{Ш}(E/K)^{\epsilon}_{l^\infty}$ follows from Lemma 5.1.7(i).

On the one hand, l^{N_1} is the maximum order of an element of $\text{Ш}(E/K)^{\epsilon}_{l^\infty}$. From Proposition 5.2.1 and the isomorphism of (5.4.4), there is an element in the Selmer group $\text{Sel}_{l^\infty}(E/K)^{\epsilon}$, and hence in the Tate-Shafarevich group $\text{Ш}(E/K)^{\epsilon}_{l^\infty}$, of order $l^{M_0-M_1}$. It follows that we have the inequality

$$M_0 - M_1 \leq N_1. \tag{5.4.11}$$

On the other hand, the Cassels pairing gives, as $l^n\delta_{M_0}(z_1) = \delta_{M_0-n}(z_1)$ if $n \leq M_0$,

$$\begin{aligned} < \delta_{M_0}(z_1), l^n d >_{\text{Cassels}} &=< l^n\delta_{M_0}(z_1), d >_{\text{Cassels}} \\ &=< \delta_{M_0-n}(z_1), d >_{\text{Cassels}} \end{aligned} \tag{5.4.12}$$

and where we have

$$l^{N_i}\delta_{M_0+N_i-n}(z_1) = \delta_{M_0-n}(z_1).$$

By the construction of the Cassels pairing as a sum of local pairings, more precisely from Proposition 4.6.1 and equation (4.6.2), we obtain from this last equation (5.4.12) that for all $0 \leq n \leq N_i - 1$ where l^{N_i} is the order of d_i

$$\begin{aligned} < \delta_{M_0}(z_1), l^n d_i >_{\text{Cassels}} &= \sum_{v\in\Sigma_K} [\delta_{M_0+N_i-n}(z_1)_v, w_{i,v}]_v \\ &= [\delta_{M_0+N_i-n}(z_1)_{y_1}, w_{i,y_1}]_{y_1} \end{aligned} \tag{5.4.13}$$

where $e_{i,v} = \partial_{l^{N_i}}(w_{i,v})$ as in (5.4.6) for all $v \in \Sigma_K$, as we have that $\delta_{M_0+N_i-n}(z_1)_v = 0$ for all places v of K not dividing z_1 (by Proposition 4.5.1(i)).

The term

$$[\delta_{M_0+N_i-n}(z_1)_{y_1}, w_{i,y_1}]_{y_1}$$

is zero if $i \geq 2$ by (5.4.10). Hence we have from (5.4.13)

$$< \delta_{M_0}(z_1), d_i >_{\text{Cassels}} = 0 \quad \text{for } i \geq 2. \tag{5.4.14}$$

Let $i = 1$. From the sum formula (5.4.13) we have

$$< \delta_{M_0}(z_1), l^n d_1 >_{\text{Cassels}} = [\delta_{M_0+N_1-n}(z_1)_{y_1}, w_{1,y_1}]_{y_1}. \tag{5.4.15}$$

By (5.4.9), the term $[\delta_{M_0+N_i}(z_1)_{y_1}, w_{1,y_1}]_{y_1}$ has order l^{N_1}. Hence by (5.4.15), the element $< \delta_{M_0}(z_1), d_1 >_{\text{Cassels}}$ of $\mathbb{Q}/\mathbb{Z}$ has order l^{N_1}. Hence the character

$$d \mapsto < \delta_{M_0}(z_1), l^n d >_{\text{Cassels}}, \quad Ш(E/K)^{\epsilon}_{l^\infty} \to \mathbb{Q}/\mathbb{Z},$$

is non-zero for all n such that $0 \leq n \leq N_1 - 1$ and more precisely

$$< \delta_{M_0}(z_1), l^n d_1 >_{\text{Cassels}}$$

is non-zero for all n such that $0 \leq n \leq N_1 - 1$.

Therefore the character

$$d \mapsto < \delta_{M_0}(z_1), d >_{\text{Cassels}}, \quad Ш(E/K)^{\epsilon}_{l^\infty} \to \mathbb{Q}/\mathbb{Z},$$

vanishes on $\prod_{i\geq 2} D_i$ (by (5.4.14)) and its restriction to D_1 generates the dual $\widehat{D_1}$. Hence the element $\delta_{M_0}(z_1)$ of $Ш(E/K)^{\epsilon}_{l^\infty}$ has order at least l^{N_1} as this is the order of the cyclic group D_1. Since $\delta_{M_0}(z_1)$ has order at most $l^{M_0-M_1}$, by definition of the cohomology class $\delta_{M_0}(z_1)$ (see Lemma 5.1.7(iii)), we obtain that

$$N_1 \leq M_0 - M_1.$$

Hence we must have from the inequality (5.4.11)

$$N_1 = M_0 - M_1. \tag{5.4.16}$$

It follows from this equality that $\delta_{M_0}(z_1)$ has order $l^{M_0-M_1}$ and therefore l^{M_1+1} does not divide P_{z_1} and hence we have

$$l^{M_1} || P_{z_1}.$$

In summary, we have shown that $\delta_{M_0}(z_1)$ has order $l^{N_1} = l^{M_0-M_1}$, $l^{M_1}||P_{z_1}$, $S_{y_1} = 0$, and the character $d \mapsto < \delta_{M_0}(z_1), d >_{\text{Cassels}}$ vanishes on $\prod_{i\geq 2} D_i$ and its restriction to D_1 generates the dual $\widehat{D_1}$.

We now proceed by induction. Suppose there are prime divisors $z_1, \ldots, z_k \in \Lambda^1(M_0 + N_1)$ such that

$$e_{i,y_j} = 0, \quad \text{for all} \quad i \neq j, \text{ and for all } j \text{ such that } 1 \leq j \leq k \tag{5.4.17}$$

and if

$$c_j = z_1 + \ldots + z_j$$

then

$$l^{M_j} || P_{c_j}, \ \delta_{M_{j-1}}(c_j) \in Ш(E/K)^{-\nu(j)}_{l^\infty}, \quad \text{for all } 1 \leq j \leq k \tag{5.4.18}$$

and the characters

$$\chi_j : d \mapsto < \delta_{M_{j-1}}(c_j), d >_{\text{Cassels}}, \ 1 \leq j \leq k$$

vanish on $\prod_{i\geq k+1} D_i$ and form a triangular basis of the dual of $\prod_{i\leq k} D_i$ such that the restriction of χ_j to the cyclic subgroup D_j is a basis for the dual $\widehat{D_j}$ for all $j = 1, \ldots, k$. Suppose further we have shown that

$$M_{j-1} - M_j = N_j, \text{ for } 1 \leq j \leq k. \tag{5.4.19}$$

and

$$\text{ord } \delta_{M_{j-1}}(c_j) = l^{N_j} \text{ for } 1 \leq j \leq k. \tag{5.4.20}$$

We have already proved the existence of the divisor $c_1 = z_1$ and these properties of the previous paragraph of e_{i,y_1}, P_{c_1}, $\delta_{M_0}(c_1)$, χ_1, $N_1 = M_0 - M_1$, including (5.4.17), (5.4.18), (5.4.19), and (5.4.20) for $k = 1$. Let $y_i \in \Sigma_K$ be the place of K above z_i for all $i = 1, \ldots, k$.

Let m be the integer already selected such that $l^m \geq \exp(Ш(E/K)_{l^\infty})$. The order of $\delta_{M_{k-1}}(c_k)$ in $Ш(E/K)_{l^\infty}$ is the same as its order as a character on D via the non-degenerate Cassels pairing. Since D is an isotropic subgroup of $Ш(E/K)_{l^\infty}$, it follows that

$$\mathbb{Z}\delta_{M_{k-1}}(c_k) \cap D = 0. \tag{5.4.21}$$

We have

$$\gamma_{M_k+N_{k+1}}(c_k) = l^{N_k - N_{k+1}} \gamma_{M_{k-1}}(c_k)$$

as $N_k = M_{k-1} - M_k$ from (5.4.19) and where $N_k - N_{k+1} \geq 0$ by the definition of the integers N_i as the invariants of $Ш(E/K)_{l^\infty}$ in decreasing order. It follows from the induction hypothesis that the cohomology class $\delta_{M_k+N_{k+1}}(c_k)$ has order $l^{N_{k+1}}$, by (5.4.20), and belongs to $Ш(E/K)_{l^\infty}^{-\nu(k)}$.

We have $l^{M_k} || P_{c_k}$ by the induction hypothesis (5.4.18). Hence by Lemma 3.4.14(i), the cohomology class $\gamma_{M_k+N_{k+1}}(c_k)$ then has the same order as its homomorphic image $\delta_{M_k+N_{k+1}}(c_k)$ namely $l^{N_{k+1}}$.

Let S be the subgroup of the Selmer group $\text{Sel}_{l^m}(E/K)$ generated by the elements e_i for all $i \neq k+1$. Let T be the subgroup of the Selmer group $\text{Sel}_{l^m}(E/K)$ generated by $\gamma_{M_k+N_{k+1}}(c_k), e_{k+1}, S$. From the isomorphism (5.4.4) and that $\mathbb{Z}\delta_{M_k+N_{k+1}}(c_k)$ has trivial intersection with D by (5.4.21), there is an equality of subgroups of the Selmer group $\text{Sel}_{l^m}(E/K)$, where the sums on the right hand side are direct,

$$T = \mathbb{Z}\gamma_{M_k+N_{k+1}}(c_k) \oplus \mathbb{Z}e_{k+1} \oplus S.$$

We may now apply Lemma 5.4.3 to S and the elements $e_{k+1}, \gamma_{M_k+N_{k+1}}(c_k)$ where we take

$$c = c_k, r = k, s = M_k + N_{k+1}, t = M_0 + N_1 - s, n = N_{k+1}. \tag{5.4.22}$$

There is then according to the lemma a prime divisor $z_{k+1} \in \Lambda^1(M_0 + N_1)$ which satisfies the following 2 conditions

$$\text{ord}[\delta_{M_k+N_{k+1}}(c_k + z_{k+1}), w_{k+1,y_{k+1}}]_{y_{k+1}} = l^{N_{k+1}} \tag{5.4.23}$$

and

$$S_{y_{k+1}} = 0 \tag{5.4.24}$$

where $y_{k+1} \in \Sigma_K$ is the prime divisor of K lying over z_{k+1} and where the subscript y_{k+1} denotes localization at y_{k+1}, and where the point $w_{k+1,y_{k+1}}$ in $(E(K_{y_{k+1}})/l^{N_{k+1}}E(K_{y_{k+1}}))^\epsilon$ is such that $\partial_{l^{N_{k+1}}}(w_{k+1,y_{k+1}}) = e_{k+1,y_{k+1}}$.

For this selection of $z_{k+1} \in \Lambda^1(M_0 + N_1)$, note that $M_k + N_{k+1} \leq M_0 + N_1$ and so $t \geq 0$, where t is the parameter of (5.4.22), because M_r, N_r are both decreasing sequences of integers and that $H^1(K, E_{l^{M_k+N_{k+1}}})$ is a subgroup of $H^1(K, E_{l^{M_0+N_1}})$ by Lemma 3.1.5.

Let c_{k+1} be the divisor which is the sum of the z_i, for $i = 1, \ldots, k+1$,

$$c_{k+1} = \sum_{j=1}^{k+1} z_j. \tag{5.4.25}$$

Let

$$\delta_{M_k}(c_{k+1}) \in Ш(E/K)_{l^\infty}^{-\nu(k+1)}$$

be the cohomology class associated to this divisor $c_{k+1} \in \Lambda^{k+1}(M_0 + N_1)$; that $\delta_{M_k}(c_{k+1})$ belongs to the Tate-Shafarevich group $Ш(E/K)_{l^\infty}^{-\nu(k+1)}$ follows from Lemma 5.1.7(i).

Then for $0 \leq n \leq N_{k+1} - 1$ by the construction of the Cassels pairing as a sum of local terms, more precisely from Proposition 4.6.1 and equation (4.6.2), we have the following sum formulae as d_i has order l^{N_i}

$$< \delta_{M_k}(c_{k+1}), l^n d_i >_{\text{Cassels}} = < \delta_{M_k-n}(c_{k+1}), d_i >_{\text{Cassels}} \tag{5.4.26}$$

$$= \sum_{v \in \Sigma_K} [\delta_{M_k-n+N_i}(c_{k+1})_v, w_{i,v}]_v = \sum_{j=1}^{k+1} [\delta_{M_k-n+N_i}(c_{k+1})_{y_j}, w_{i,y_j}]_{y_j}.$$

because $\delta_{M_k-n+N_i}(c_{k+1})_v = 0$ for all $v \in \Sigma_K$ not dividing an element of $\text{Supp}(c_{k+1})$ by Proposition 4.5.1(i). Here $w_{i,y_j} \in {}_{l^{N_i}}E(K_{y_j})$ is an element already chosen (see (5.4.6)) such that

$$\partial_{l^{N_i}}(w_{i,y_j}) = e_{i,y_j}.$$

We have the following sum for the Cassels pairing, obtained from those of (5.4.26),

$$< \delta_{M_k}(c_{k+1}), l^n d_i >_{\text{Cassels}} = \sum_{j=1}^{k+1} [\delta_{M_k-n+N_i}(c_{k+1})_{y_j}, w_{i,y_j}]_{y_j}. \tag{5.4.27}$$

We have $w_{i,y_j} = 0$ for all $i \neq j$ and all j such that $1 \leq j \leq k+1$ by (5.4.17), for $j \leq k$, and by (5.4.24), for $j = k+1$. It follows that for $i \geq k+1$ all terms $[\delta_{M_k - n + N_i}(c_{k+1})_{y_j}, w_{i,y_j}]_{y_j}$ of this sum (5.4.27) are zero except the last $[\delta_{M_k - n + N_i}(c_{k+1})_{y_{k+1}}, w_{i,y_{k+1}}]_{y_{k+1}}$ and the entire sum is zero for $i > k+1$.

By (5.4.23) the local term

$$[\delta_{M_k + N_{k+1} - n}(c_{k+1})_{y_{k+1}}, w_{k+1,y_{k+1}}]_{y_{k+1}}$$

is non-zero for all integers n such that $0 \leq n \leq N_{k+1} - 1$.

Therefore the character, by (5.4.27),

$$\chi_{k+1} : \ d \mapsto < \delta_{M_k}(c_{k+1}), d >_{\text{Cassels}}$$

vanishes on $\prod_{i>k+1} D_i$ and its restriction to D_{k+1} generates $\widehat{D}_{k+1}$, as D_{k+1} has order $l^{N_{k+1}}$ by definition. Hence χ_{k+1} extends the triangular basis $\chi_1, \ldots, \chi_k$ to generate $\prod_{i \leq k+1} \widehat{D}_i$ and $\delta_{M_k}(c_{k+1})$ has order at least $l^{N_{k+1}}$. Since it has order at most $l^{M_k - M_{k+1}}$, by definition of the cohomology class $\delta_{M_k}(c_{k+1})$ (see Lemma 5.1.7(iii)), we conclude that

$$N_{k+1} \leq M_k - M_{k+1} \tag{5.4.28}$$

and also

$$l^{N_{k+1}} \leq \ \text{ord}\ \delta_{M_k}(c_{k+1}) \ \leq l^{M_k - M_{k+1}}. \tag{5.4.29}$$

Let C_k be the subgroup of the Selmer eigencomponent $\text{Sel}_{l^m}(E/K)^{-\nu(k+1)}$ given by

$$C_k = \left(\gamma_{M_0+N_1}(0), e_1, \ldots, e_k, \gamma_{M_0}(c_1), \ldots, \gamma_{M_{k-1}}(c_k)\right)^{-\nu(k+1)}.$$

If k is even then C_k is generated by $e_1, \gamma_{M_0}(c_1), e_3, \gamma_{M_2}(c_3), \ldots, e_{k-1}, \gamma_{M_{k-2}}(c_{k-1})$ of which there are k in number. If k is odd then C_k is generated by $\gamma_{M_0+N_1}(0), e_2, \gamma_{M_1}(c_2), e_4, \gamma_{M_3}(c_4), \ldots, e_{k-1}, \gamma_{M_{k-2}}(c_{k-1})$ of which there are k in number. We then have for all integers k

$$\text{rank}(C_k) \leq k. \tag{5.4.30}$$

The elements $e_1, \ldots, e_k$ generate a subgroup of $\text{Sel}_{l^m}(E/K)$ isomorphic to $\prod_{i \leq k} D_i$ by the decomposition (5.4.4) of the Selmer group. Furthermore, the elements $\gamma_{M_0}(c_1), \ldots, \gamma_{M_{k-1}}(c_k)$ generate a subgroup of $\text{Sel}_{l^m}(E/K)$ isomorphic to the dual of $\prod_{i \leq k} D_i$ as $\gamma_{M_{i-1}}(c_i)$ has the same order as $\delta_{M_{i-1}}(c_i)$ for all $i = 1, \ldots, k$ (see (5.4.18), (5.4.19), (5.4.20), and Proposition 3.4.14(i)). In the decomposition (5.4.4) of $\text{Sel}_{l^m}(E/K)$, take S to be the subgroup ${}_{l^m}E(K) \oplus_{1 \leq i \leq k} \prod D_i \oplus \bigoplus_{1 \leq i \leq k} \mathbb{Z}\gamma_{M_{i-1}}(c_i)$. Then we have $C_k = S^{-\nu(k+1)}$.

If a finite abelian group G is a direct product $G_1 \times G_2$ of 2 subgroups and $g \in G$ is such that $\mathbb{Z}g \cap G_1 = 0$ then the order of the element g is at most the exponent of G_2. Then by the previous remark where we take $G = \mathrm{Sel}_{l^m}(E/K)^{-\nu(k+1)}$ and $G_1 = C_k$, we have that $l^{N_{k+1}}$ is the maximum order of an element $c \in \mathrm{Sel}_{l^m}(E/K)^{-\nu(k+1)}$ if

$$\mathbb{Z}c \cap C_k = 0$$

by the decomposition (5.4.4) of the Selmer group $\mathrm{Sel}_{l^m}(E/K)$.

On the other hand, by Proposition 5.2.1 and (5.4.30) applied to subgroup C_k of the Selmer group there is an element in $\mathrm{Sel}_{l^m}(E/K)^{-\nu(k+1)}$ of order $l^{M_k - M_{k+1}}$ satisfying $\mathbb{Z}c \cap C_k = 0$. Hence we have

$$M_k - M_{k+1} \leq N_{k+1}$$

and so by (5.4.28) we have

$$N_{k+1} = M_k - M_{k+1}. \tag{5.4.31}$$

It follows from this equality and (5.4.29) that

$$\text{ord } \delta_{M_k}(c_{k+1}) = l^{M_k - M_{k+1}}. \tag{5.4.32}$$

Therefore $l^{M_{k+1}+1}$ does not divide $P_{c_{k+1}}$ and $l^{M_{k+1}} | P_{c_{k+1}}$ and hence we have

$$l^{M_{k+1}} || P_{c_{k+1}}. \tag{5.4.33}$$

The properties (5.4.31), (5.4.32), and (5.4.33) complete the proof of the induction step and this proves the Theorems 4.1.9 and 4.1.13. □

5.5. Proof of Theorems 1.1.1. and 4.1.14

Proof of Theorem 1.1.1. From Proposition 2.2.5, we have as $l \in \mathcal{P}$ is an odd prime number that $Ш(E/F)_{l^\infty} \cong Ш(E/K)_{l^\infty}^{+1}$. The Theorem now follows immediately from Theorem 4.1.9. □

Proof of Theorem 4.1.14. As P_0 has infinite order, by Lemma 5.1.2 and [1, Chap. 7, Theorem 7.6.5] we have that M_0 is finite, the group $\mathbb{Z}P_0$ has finite index in $E(K)$, and the highest power of l dividing the index $[E(K) : \mathbb{Z}P_0]$ is equal to

$$l^{M_0} = |(E(K)/\mathbb{Z}P_0)_{l^\infty}|.$$

By Theorem 4.1.9, we have that

$$\left| Ш(E/K)_{l^\infty} \right| = \prod_{i \geq 0} l^{2(M_i - M_{i+1})} = l^{2(M_0 - M_\infty)}$$

where

$$M_\infty = \min_{i \in \mathbf{N}} M_i$$

and where this minimum exists as the M_i form a decreasing sequence of non-negative integers by Lemma 5.1.4. □

5.6. Generators of Tate-Shafarevich groups

(5.6.1) The notation (5.1.1) of the section §5.1 holds also for this section. We further denote by

Σ_K the set of all places of the global field K;
$[,]_v : {}_{l^m}E(K_v) \times H^1(K_v, E)_{l^m} \to \mathbb{Z}/l^m\mathbb{Z}$ the non-degenerate local pairing at the place v of K induced by the cup product, as in Theorem 2.1.6.

5.6.2. Theorem. *Let l be a prime number belonging to $\mathcal{P}$; assume that l is coprime to the order of* $\mathrm{Pic}(A)$. *Suppose that P_0 has infinite order in $E(K)$ and let a be an integer such that $a \geq 2M_0$. Then we have*
(1) *the classes $\delta_{M_0}(c)$, for all $c \in \Lambda^1(a)$, generate $Ш(E/K)^\epsilon_{l^\infty}$;*
(2) *the classes $\delta_{M_1}(c)$, for all $c \in \Lambda^2(a)$, generate $Ш(E/K)^{-\epsilon}_{l^\infty}$.*

5.6.3. Theorem. *Under the hypotheses of Theorem 5.6.2, the classes $\delta_{M_r}(c)$, for all $c \in \Lambda^r(a)$, generate $Ш(E/F)_{l^\infty}$ where $r = (3-\epsilon)/2$.*

Proof of Theorem 5.6.3. This evidently follows from Theorem 5.6.2 and Proposition 2.2.5. □

Proof of Theorem 5.6.2. By Theorem 4.1.9 above or [1, Chap. 7, Theorem 7.6.5], the group $Ш(E/K)_{l^\infty}$ is finite. By Lemmas 5.1.2 and 5.1.4, the quantities $M_0, M_1, M_2, \ldots$ are all finite and form a decreasing sequence of non-negative integers. We fix an integer $a \geq 2M_0$. The classes $\delta_{M_0}(z)$, for $z \in \Lambda^1(a)$, and the classes $\delta_{M_1}(z_1 + z_2)$, for $z_1 + z_2 \in \Lambda^2(a)$, belong to $Ш(E/K)_{l^\infty}$ by Lemma 5.1.7(i). We now prove separately the two parts of the theorem.

(1) Suppose that $d \in Ш(E/K)^\epsilon_{l^\infty}$ has order exactly l^M for some $M > 0$ and is in the ϵ-eigenspace of $Ш(E/K)_{l^\infty}$. By Theorem 4.1.15 there is an isomorphism of ϵ-components

$$\mathrm{Sel}_{l^\infty}(E/K)^\epsilon \cong Ш(E/K)^\epsilon_{l^\infty} \tag{5.6.4}$$

induced from the natural surjection of the Selmer group onto the Tate-Shafarevich group.

By Theorem 4.1.9, we have $M \leq M_0$. By the isomorphism (5.6.4), we may lift d to an element of the Selmer group $e \in \mathrm{Sel}_{l^\infty}(E/K)^\epsilon$ of order l^M. The cohomology class $\gamma_{M_0+M}(0)$ belongs to the $-\epsilon$-eigenspace $\mathrm{Sel}_{l^\infty}(E/K)^{-\epsilon}$ of the Selmer group and has order l^M by Lemma 5.1.2.

We may apply Lemma 5.4.3 to the elements e, $\gamma_{M_0+M}(0)$ and the subgroup $S = 0$ where we take the parameters of the lemma to be

$$c = 0, r = 0, s = M_0 + M, t = a - (M_0 + M), n = M. \tag{5.6.5}$$

Note that $t \geq 0$ as $a \geq 2M_0 \geq M_0 + M$. There is according to the lemma a prime divisor $z \in \Lambda^1(a)$ which satisfies the following condition

$$\mathrm{ord}[\delta_{M_0+M}(z)_y, w_y]_y = l^M \tag{5.6.6}$$

where $y \in \Sigma_K$ is the prime divisor of K lying over z and where the subscript y denotes localization at y, and where the point w_y in $(E(K_y)/l^M E(K_y))^\epsilon$ is such that $\partial_{l^M}(w_y) = e_y$. Here in (5.6.6), $\delta_{M_0+M}(z)$ is the cohomology class in $H^1(K, E)_{l^{M_0+M}}$ associated to z.

Let $\delta_{M_0}(z)$ be the cohomology class of $Ш(E/K)^\epsilon_{l^\infty}$ associated to z, where this class belongs to the Tate-Shafarevich group by Lemma 5.1.7.

The Cassels pairing gives, as $l^u \delta_{M_0}(z) = \delta_{M_0-u}(z)$ if $u \leq M_0$,

$$< \delta_{M_0}(z), l^u d >_{\mathrm{Cassels}} = < \delta_{M_0-u}(z), d >_{\mathrm{Cassels}} . \tag{5.6.7}$$

As d has order l^M and that we have

$$l^M \delta_{M_0+M-u}(z) = \delta_{M_0-u}(z),$$

by the construction of the Cassels pairing in terms of local Tate pairings (Proposition 4.6.1 and equation (4.6.2)) we obtain from this last equation (5.6.7)

$$< \delta_{M_0}(z), l^u d >_{\mathrm{Cassels}} = [\delta_{M_0+M-u}(z)_y, w_y]_y$$

where as above

$$e_y = \partial_{l^M}(w_y),$$

and where $w_y \in {}_{l^M}E(K_y)$ as we have that $\delta_{M_0+M-u}(z)_v = 0$ for all places v of K not dividing z by Proposition 4.5.1(i).

By (5.6.6) the element given by the Tate pairing

$$[\delta_{M_0+M-u}(z)_y, w_y]_y$$

of $\mathbb{Q}/\mathbb{Z}$ is non-zero for all integers u such that $1 \leq u \leq M - 1$ and hence

$$< \delta_{M_0}(z), l^u d >_{\mathrm{Cassels}}$$

is non-zero for all integers u such that $1 \leq u \leq M-1$. We obtain that the character, where $z \in \Lambda^1(a)$,

$$f \mapsto < \delta_{M_0}(z), f >_{\text{Cassels}}, \ \text{Ш}(E/K)_{l^\infty} \to \mathbb{Q}/\mathbb{Z},$$

of $\text{Ш}(E/K)_{l^\infty}$ when restricted to $\mathbb{Z}d$ generates the dual $\widehat{\mathbb{Z}d}$ of this subgroup $\mathbb{Z}d$.

As d is any element of the abelian group $\text{Ш}(E/K)^\epsilon_{l^\infty}$, the non-degeneracy of Cassels pairing implies that the classes $\{\delta_{M_0}(z), z \in \Lambda^1(a)\}$ generate the ϵ-eigenspace $\text{Ш}(E/K)^\epsilon_{l^\infty}$ which proves the part of the theorem for $\text{Ш}(E/K)^\epsilon_{l^\infty}$.

(2) Suppose that $f \in \text{Ш}(E/K)^{-\epsilon}_{l^\infty}$ has order exactly $l^{M'}$ and is in the $-\epsilon$-eigenspace of $\text{Ш}(E/K)_{l^\infty}$. We have $M' \leq M_0$ by Theorem 4.1.9.

We have from Theorem 4.1.15, an isomorphism compatible with the τ-eigenspaces

$$\text{Sel}_{l^m}(E/K) \cong {}_{l^m}E(K) \oplus \text{Ш}(E/K)_{l^\infty} \ \text{ for all } m \geq N_1$$

from which as stated in this theorem we obtain the isomorphism, taking $m = a \geq 2M_0 \geq N_1$,

$$\Delta_a : \text{Sel}_{l^a}(E/K)^{-\epsilon} \cong \mathbb{Z}/l^a\mathbb{Z} \oplus \text{Ш}(E/K)^{-\epsilon}_{l^\infty}.$$

The projection of $\text{Sel}_{l^a}(E/K)$ onto the second factor $\text{Ш}(E/K)_{l^\infty}$ of this direct sum is the natural surjection of the Selmer group onto the Tate-Shafarevich group.

Lift f to $g \in \text{Sel}_{l^a}(E/K)^{-\epsilon}$ via this isomorphism Δ_a for the integer a and where g has order $l^{M'}$ and has zero component in the first term $\mathbb{Z}/l^a\mathbb{Z}$ of this decomposition of the Selmer group.

By Theorem 4.1.9 or alternatively Proposition 5.2.1, there is an element $d \in \text{Ш}(E/K)^\epsilon_{l^\infty}$ of order exactly $l^{M_0-M_1}$. Lift, via the isomorphism (5.6.4), d to an element of the Selmer group $e \in \text{Sel}_{l^a}(E/K)^\epsilon$ which is also of order $l^{M_0-M_1}$.

The cohomology class $\gamma_{2M_0-M_1}(0)$ belongs to the $-\epsilon$-eigenspace $\text{Sel}_{l^a}(E/K)^{-\epsilon}$ of the Selmer group and has order $l^{M_0-M_1}$ by Lemma 5.1.2.

Let T be the subgroup of $\text{Sel}_{l^a}(E/K)$ generated by the three elements $\gamma_{2M_0-M_1}(0), e$ and g. As the two elements $\gamma_{2M_0-M_1}(0), g$ belong to the different components of $\text{Sel}_{l^a}(E/K)^{-\epsilon}$ under the isomorphism Δ_a and as e belongs to a different eigenspace $\text{Sel}_{l^a}(E/K)^\epsilon$, the group T is the direct sum of the subgroups generated by these three elements, that is to say we have

$$T \cong \mathbb{Z}\gamma_{2M_0-M_1}(0) \oplus \mathbb{Z}e \oplus \mathbb{Z}g.$$

We may apply Lemma 5.4.3 to the subgroup $S = \mathbb{Z}g$ and the elements $\gamma_{2M_0-M_1}(0)$ and e of the Selmer group; we take the parameters of the lemma to be

$$c = 0, r = 0, s = 2M_0 - M_1, t = a - (2M_0 - M_1), n = M_0 - M_1. \tag{5.6.8}$$

Note that $t \geq 0$ as $a \geq 2M_0$. There is according to the lemma a prime divisor $z_1 \in \Lambda^1(a)$ which satisfies the following 2 conditions

$$\text{ord}[\delta_{2M_0-M_1}(z_1)_{y_1}, w_{y_1}]_{y_1} = l^{M_0-M_1} \tag{5.6.9}$$

$$(\mathbb{Z}g)_{y_1} = 0 \tag{5.6.10}$$

where $y_1 \in \Sigma_K$ is the prime divisor of K lying over z_1, and where the point w_{y_1} in $(E(K_{y_1})/l^{M_0-M_1}E(K_{y_1}))^\epsilon$ is such that $\partial_{l^{M_0-M_1}}(w_{y_1}) = e_{y_1}$. Here in (5.6.9), $\delta_{2M_0-M_1}(z_1)$ is the cohomology class in $H^1(K,E)_{l^{2M_0-M_1}}$ associated to z_1.

Let

$$\delta_{M_0}(z_1) \in \text{Ш}(E/K)^\epsilon_{l^\infty}$$

be the cohomology class associated to this prime divisor $z_1 \in \Lambda^1(a)$; that $\delta_{M_0}(z_1)$ belongs to $\text{Ш}(E/K)^\epsilon_{l^\infty}$ follows from Lemma 5.1.7(i).

The Cassels pairing gives, as $l^u\delta_{M_0}(z_1) = \delta_{M_0-u}(z_1)$ if $u \leq M_0$,

$$< \delta_{M_0}(z_1), l^u d >_{\text{Cassels}} =< l^u\delta_{M_0}(z_1), d >_{\text{Cassels}} \tag{5.6.11}$$

$$=< \delta_{M_0-u}(z_1), d >_{\text{Cassels}}$$

where

$$l^{M_0-M_1}\delta_{2M_0-M_1-u}(z_1) = \delta_{M_0-u}(z_1).$$

By the construction of the Cassels pairing as a sum of local terms, more precisely from Proposition 4.6.1 and equation (4.6.2), we obtain from this last equation (5.6.11) that for all $0 \leq u \leq M_0 - M_1 - 1$ where $M_0 - M_1$ is the order of d

$$< \delta_{M_0}(z_1), l^u d >_{\text{Cassels}} = \sum_{v\in\Sigma_K} [\delta_{2M_0-M_1-u}(z_1)_v, w_v]_v \tag{5.6.12}$$

$$= [\delta_{2M_0-M_1-u}(z_1)_{y_1}, w_{y_1}]_{y_1}$$

where $e_v = \partial_{l^{M_0-M_1}}(w_v)$ for all $v \in \Sigma_K$, and where we have that $\delta_{2M_0-M_1-u}(z_1)_v = 0$ for all places v of K not dividing z_1 (by Proposition 4.5.1(i)).

From the sum formula (5.6.12) we have

$$< \delta_{M_0}(z_1), l^u d >_{\text{Cassels}} = [\delta_{2M_0-M_1-u}(z_1)_{y_1}, w_{y_1}]_{y_1}.$$

By (5.6.9), $[\delta_{2M_0-M_1}(z_1)_{y_1}, w_{y_1}]_{y_1}$ has order $l^{M_0-M_1}$.

Hence the map

$$s \mapsto < \delta_{M_0}(z_1), l^u s >_{\text{Cassels}}, \quad \text{Ш}(E/K)^\epsilon_{l^\infty} \to \mathbb{Q}/\mathbb{Z},$$

is non-zero for all u such that $0 \leq u \leq M_0 - M_1 - 1$ and more precisely

$$< \delta_{M_0}(z_1), l^u d >_{\text{Cassels}}$$

is non-zero for all u such that $0 \leq u \leq M_0 - M_1 - 1$.

Therefore the character

$$s \mapsto < \delta_{M_0}(z_1), s >_{\text{Cassels}}, \quad Ш(E/K)^{\epsilon}_{l^\infty} \to \mathbb{Q}/\mathbb{Z},$$

is such that its restriction to $\mathbb{Z}d$ generates the dual $\widehat{\mathbb{Z}d}$. Hence $\delta_{M_0}(z_1)$ has order at least $l^{M_0-M_1}$ as this is the order of the cyclic group $\mathbb{Z}d$. Since $\delta_{M_0}(z_1)$ has order at most $l^{M_0-M_1}$, by definition of the cohomology class $\delta_{M_0}(z_1)$ (see Lemma 5.1.7(iii)), we obtain that $\delta_{M_0}(z_1)$ has order given by

$$\text{ord } \delta_{M_0}(z_1) = l^{M_0-M_1}.$$

We have $l^{M_1}|P_{z_1}$ by definition of M_1. It follows that $\gamma_{M_0}(z_1)$ has order $\leq l^{M_0-M_1}$ by Proposition 3.4.14(i). As $\delta_{M_0}(z_1)$ has order $l^{M_0-M_1}$ by the previous paragraph and $\delta_{M_0}(z_1)$ is a homomorphic image of $\gamma_{M_0}(z_1)$ we must have that $\gamma_{M_0}(z_1)$ has order exactly given by

$$\text{ord } \gamma_{M_0}(z_1) = l^{M_0-M_1}.$$

Therefore we have by Proposition 3.4.14(i)

$$l^{M_1}||P_{z_1}.$$

We may apply Lemma 5.4.3 to the subgroup $S = 0$ and the two elements $\gamma_{M_1+M'}(z_1)$ and g of the Selmer group which both have order $l^{M'}$; we take the parameters of the lemma to be

$$c = z_1, r = 1, s = M_1 + M', t = a - (M_1 + M'), n = M'. \tag{5.6.13}$$

Note that $t \geq 0$ as $a \geq 2M_0$ and $M', M_1 \leq M_0$. There is according to the lemma a prime divisor $z_2 \in \Lambda^1(a)$ which satisfies the following condition

$$\text{ord}[\delta_{M_1+M'}(z_1 + z_2)_{y_2}, x_{y_2}]_{y_2} = l^{M'} \tag{5.6.14}$$

where $y_2 \in \Sigma_K$ is the prime divisor of K lying over z_2, and where the point x_{y_2} in $(E(K_{y_2})/l^{M'}E(K_{y_2}))^{\epsilon}$ is such that $\partial_{l^{M'}}(x_{y_2}) = e_{y_2}$. Here in (5.6.14), $\delta_{M_1+M'}(z_1 + z_2)$ is the cohomology class in $H^1(K, E)_{l^{M_1+M'}}$ associated to $z_1 + z_2$.

Let

$$\delta_{M_1}(z_1 + z_2)$$

be the cohomology class of $Ш(E/K)^{-\epsilon}_{l^\infty}$ associated to the divisor $z_1 + z_2 \in \Lambda^2(a)$, where this class belongs to the Tate-Shafarevich group by Lemma 5.1.7(i).

Then the Cassels pairing gives, as $l^u\delta_{M_1}(z_1 + z_2) = \delta_{M_1-u}(z_1 + z_2)$ if $u \leq M_1$,

$$< \delta_{M_1}(z_1 + z_2), l^u f >_{\text{Cassels}} = < \delta_{M_1-u}(z_1 + z_2), f >_{\text{Cassels}} . \tag{5.6.15}$$

As f has order $l^{M'}$ and that we have

$$l^{M'}\delta_{M_1+M'-u}(z_1+z_2)=\delta_{M_1-u}(z_1+z_2),$$

by the construction of the Cassels pairing (Proposition 4.6.1 and equation (4.6.2)) we obtain from this last equation (5.6.15)

$$(5.6.16)\qquad <\delta_{M_1}(z_1+z_2), l^u f>_{\text{Cassels}}=\sum_{z\in\ \text{Supp}(z_1+z_2)}[\delta_{M_1+M'-u}(z_1+z_2)_y, x_y]_y$$

where

$$(5.6.17)\qquad g_y=\partial_{l^{M'}}(x_y),$$

and where

$$x_y\in({}_{l^{M'}}E(K_y))^{-\epsilon}$$

and because we have that $\delta_{M_1+M-u}(z_1+z_2)_v=0$ for all places v of K not dividing z_1+z_2 by Proposition 4.5.1(i). Here in (5.6.16) and (5.6.17), z runs over the places z_1, z_2 and $y\in\Sigma_K$ runs over the place of K above z, that is to say above z_1, z_2.

The first term $[\delta_{M_1+M'-u}(z_1+z_2)_{y_1}, x_{y_1}]_{y_1}$ in the sum (5.6.16) is zero by (5.6.10) and (5.6.17).

By (5.6.14), the second term, an element of $\mathbb{Q}/\mathbb{Z}$,

$$[\delta_{M_1+M'-u}(z_1+z_2)_{y_2}, x_{y_2}]_{y_2}$$

is non-zero for $0\leq u\leq M'-1$ and hence $\delta_{M_1}(z_1+z_2)$ has order at least $l^{M'}$. We obtain from this and (5.6.16) that the Cassels pairing

$$<\delta_{M_1}(z_1+z_2), l^u f>_{\text{Cassels}}=[\delta_{M_1+M'-u}(z_1+z_2)_{y_1}, x_{y_1}]_{y_1}$$

is non-zero for all u such that $0\leq u\leq M'-1$. Hence this character

$$s\mapsto <\delta_{M_1}(z_1+z_2), s>_{\text{Cassels}},\quad Ш(E/K)^{-\epsilon}_{l^\infty}\to\mathbb{Q}/\mathbb{Z}$$

has restriction to $\mathbb{Z}f$ which generates the dual of this subgroup $\mathbb{Z}f$ of order $l^{M'}$.

As f is any element of the abelian group $Ш(E/K)^{-\epsilon}_{l^\infty}$, the non-degeneracy of the Cassels pairing on $Ш(E/K)^{-\epsilon}_{l^\infty}$ implies that the classes $\delta_{M_1}(z_1+z_2)$, for divisors $z_1+z_2\in\Lambda^2(a)$, generate the $-\epsilon$ eigenspace $Ш(E/K)^{-\epsilon}_{l^\infty}$ which proves the part of the theorem for $Ш(E/K)^{-\epsilon}_{l^\infty}$. □

References

[1] Brown M.L., Heegner Modules and Elliptic Curves. Springer Lecture Notes in Mathematics No. 1849. Springer Verlag, Berlin-Heidelberg-New York, 2004.
[2] Gross B.H., Kolyvagin's work on modular elliptic curves. In: L-functions and Arithmetic (eds. J.H. Coates and M.J. Taylor) Cambridge University Press 1990, pp. 235-256.
[3] Gross B.H., Zagier D., Heegner points and derivatives of L-series. Invent. Math. **84**, 225-320 (1986)
[4] Kolyvagin V.A., Euler systems. In: The Grothendieck Festschrift (Vol. II). P. Cartier et al., eds., pp. 435-483. Progress in Mathematics 8. Birkhäuser, Boston 1990
[5] Kolyvagin V.A., Finiteness of $E(\mathbb{Q})$ and $Ш(E/\mathbb{Q})$ for a class of Weil curves. Math. USSR Vol. **32** (1989) No. 3.
[6] Kolyvagin V.A., On the structure of Shafarevich-Tate groups, Proceedings of USA-USSR Symposium on Algebraic Geometry, Chicago 1989. Lecture Notes in Mathematics No. 1479, Springer Verlag, Berlin-Heidelberg-New York 1991
[7] Kolyvagin V.A., On the structure of Selmer groups, Math. Annalen **291**, 253-259 (1991)
[8] Mccallum W.G., Kolyvagin's work on Shafarevich-Tate groups. In: L-functions and Arithmetic (eds. J.H. Coates and M.J. Taylor) Cambridge University Press 1990, pp. 295-316.
[9] Milne J.S., Arithmetic Duality Theorems, Perspectives in Mathematics, Academic Press, Boston 1986
[10] Raynaud, M., Caractéristique d'Euler-Poincaré d'un faisceau et cohomologie des variétés abéliennes. In: Dix Exposés sur la Cohomologie des Schémas. Ed. A. Grothendieck, pp. 12-30. North Holland Publ. Comp. Amsterdam 1967.